

Figure S1. Weekly ratings of pain on the average relative to the mean and maximum daily ratings for each 7-day reporting interval.




Figure legend

■ Weekly rating ■ Mean of 7 days ■ Maximum of 7 days

Figure S2. Weekly ratings of pain at its worst relative to the mean and maximum daily ratings for each 7-day reporting interval.




Figure legend

■ Weekly rating ■ Mean of 7 days ■ Maximum of 7 days

Table S1. Change in pain from baseline to week 12, by rating type and including only observations from participants who provided evaluable ratings at both time points.

Construct	Change in rating		Effect size ^a
	Mean (SE)	<i>T</i> -statistic (Significance)	
Pain on the average			
Weekly rating	-1.86 (0.22)	-8.33 (<0.001)	-1.16
Mean of 7 days	-1.91 (0.19)	-9.79 (<0.001)	-1.32
Maximum of 7 days	-2.09 (0.23)	-9.10 (<0.001)	-1.24
Pain at its worst			
Weekly rating	-2.21 (0.33)	-6.61 (<0.001)	-1.03
Mean of 7 days	-2.34 (0.24)	-9.71 (<0.001)	-1.41
Maximum of 7 days	-2.43 (0.29)	-8.34 (<0.001)	-1.27

SE: standard error.

^a Calculated as Cohen's *d*. *T*-statistics and effect sizes with the largest absolute value are in bold.

Table S2. Change in pain from baseline to week 12 of treatment, by rating type and stratified by randomized treatment assignment.

Construct	Randomized to cabozantinib			Randomized to mitoxantrone-prednisone		
	Change in rating Mean (SE)	T-statistic (Significance)	Effect size ^a	Change in rating Mean (SE)	T-statistic (Significance)	Effect size ^a
Pain on the average						
Weekly rating	-2.18 (0.29)	-7.46 (<0.001)	-1.34	-1.78 (0.33)	-5.40 (<0.001)	-1.09
Mean of 7 days	-2.13 (0.26)	-8.31 (<0.001)	-1.50	-1.81 (0.29)	-6.18 (<0.001)	-1.23
Maximum of 7 days	-2.37 (0.33)	-7.25 (<0.001)	-1.35	-2.04 (0.35)	-5.88 (<0.001)	-1.17
Pain at its worst						
Weekly rating	-2.46 (0.43)	-5.76 (<0.001)	-1.26	-1.91 (0.45)	-4.21(<0.001)	-0.89
Mean of 7 days	-2.62 (0.32)	-8.06 (<0.001)	-1.76	-1.98 (0.35)	-5.70 (<0.001)	-1.17
Maximum of 7 days	-2.71 (0.40)	-6.73 (<0.001)	-1.51	-1.94 (0.41)	-4.79 (<0.001)	-1.02

SE: standard error.

^a Calculated as Cohen's *d*. T-statistics and effect sizes with the largest absolute value are in bold.

Table S3. Change in pain from baseline to week 12 of treatment, by rating type and stratified by baseline pain ratings.

Construct	Provided a baseline pain rating ≤ 6			Provided a baseline pain rating > 6		
	Change in rating	T-statistic	Effect size ^a	Change in rating	T-statistic	Effect size ^a
	Mean (SE)	(Significance)		Mean (SE)	(Significance)	
Pain on the average						
Weekly rating	-1.88 (0.22)	-8.69 (<0.001)	1.32	-3.27 (0.97)	-3.37 (=0.007)	2.48
Mean of 7 days	-1.80 (0.19)	-9.56 (<0.001)	1.44	-3.29 (0.88)	-3.74 (=0.002)	2.19
Maximum of 7 days	-1.88 (0.24)	-7.77 (<0.001)	1.34	-3.21 (0.54)	-6.00 (<0.001)	2.56
Pain at its worst						
Weekly rating	-1.09 (0.38)	-2.86 (=0.006)	0.70	-3.20 (0.43)	-7.46 (<0.001)	1.75
Mean of 7 days	-1.74 (0.27)	-6.39 (<0.001)	1.44	-3.01 (0.39)	-7.67 (<0.001)	1.97
Maximum of 7 days	-1.28 (0.34)	-3.79 (=0.001)	1.06	-3.00 (0.38)	-7.89 (<0.001)	1.69

SE: standard error.

^a Calculated as Cohen's *d*. T-statistics and effect sizes with the largest absolute value are in bold.