Supporting Information

Development of New Benzylpiperazine Derivatives as σ_1 Receptor Ligands with *in Vivo* Antinociceptive and Anti-Allodynic Effects

Giuseppe Romeo,[†] Federica Bonanno,[†] Lisa L. Wilson,[‡] Emanuela Arena,[†] Maria N. Modica,[†] Valeria Pittalà,[†] Loredana Salerno,[†] Orazio Prezzavento,[†] Jay P. McLaughlin,[‡] and Sebastiano Intagliata^{*,†}

[†]Department of Drug and Health Sciences, University of Catania, viale A. Doria 6, 95125 Catania, Italy

[‡]Department of Pharmacodynamics, College of Pharmacy, University of Florida, Gainesville, Florida 32610, United States

*Phone: +39-095-738-4053. Email: s.intagliata@unict.it.

Contents:

 Figures S1–S18: ¹H NMR and ¹³C NMR spectra of compounds 13–16, 20–22, 23, and 24
 p. S2–S10



Figure S1. ¹H NMR spectrum of compound 13



Figure S2. ¹³C NMR spectrum of compound 13



Figure S3. ¹H NMR spectrum of compound 14



Figure S4. ¹³C NMR spectrum of compound 14



Figure S5. ¹H NMR spectrum of compound 15 (free base)



Figure S6. ¹H NMR spectrum of compound 15 (oxalate salt)



Figure S7. ¹³C NMR spectrum of compound 15 (oxalate salt).



Figure S8. ¹H NMR spectrum of compound 16





Figure S10. ¹H NMR spectrum of compound 20



Figure S11. ¹³C NMR spectrum of compound 20



Figure S12. ¹H NMR spectrum of compound 21



Figure S13. ¹³C NMR spectrum of compound 21



Figure S14. ¹H NMR spectrum of compound 22



Figure S15. ¹³C NMR spectrum of compound 22



Figure S16. ¹H NMR spectrum of compound 23



Figure S17. ¹H NMR spectrum of compound 24



Figure S18. ¹³C NMR spectrum of compound 24