

Table S2: Propensity score matched models of the relationship between mortality outcomes and dominant sedation strategy prior to first ICDSC assessment

Outcome	Dominant Sedation Strategy	Mortality, n (%)	Propensity score-matched OR (95% CI)²
ICU Mortality	Propofol	94 (6.7)	1.00 (reference)
	Fentanyl	104 (9.7)	1.50 (1.07-2.12)
	Midazolam	39 (11.0)	1.20 (0.74-1.97)
Hospital Mortality	Propofol	157 (11.1)	1.00 (reference)
	Fentanyl	166 (15.5)	1.27 (0.97-1.67)
	Midazolam	59 (16.6)	1.14 (0.76-1.70)
Died within 30 days of ICU admission	Propofol	148 (10.5)	1.00 (reference)
	Fentanyl	148 (13.8)	1.35 (1.02-1.79)
	Midazolam	50 (14.0)	1.02 (0.67-1.57)
Died within 1 year of ICU admission	Propofol	268 (19.0)	1.00 (reference)
	Fentanyl	248 (23.2)	1.13 (0.90-1.43)
	Midazolam	91 (25.6)	1.01 (0.72-1.42)
Died within 1.5 years of ICU admission	Propofol	308 (21.8)	1.00 (reference)
	Fentanyl	276 (25.8)	1.12 (0.90-1.40)
	Midazolam	109 (30.6)	1.01 (0.74-1.39)