

**Supplementary Table. Neuropsychological Test Scores (Baseline) by Treatment Group**

	<b>Total Sample (n = 101)</b>		<b>Occupational Therapy (n = 50)</b>		<b>Diabetes Education (n = 51)</b>	
Wide Range Achievement Test <sup>1</sup>						
(mean, SD)	24.0	7.5	23.7	7.7	24.3	7.5
Mini Mental State Exam <sup>2</sup>						
(mean, SD)	25.4	2.6	25.7	2.8	25.2	2.4
Digit Symbol Substitution <sup>3</sup>						
(mean, SD)	33.0	13.4	35.0	13.6	31.0	13.2
Trail Making Test-Part A <sup>4</sup>						
(mean, SD)	67.5	32.4	67.1	35.0	67.9	30.1
Trail Making Test-Part B <sup>5</sup>						
(mean, SD)	217.8	80.5	215.3	79.2	220.1	82.5
Logical Memory Immediate <sup>6</sup>						
(mean, SD)	8.6	2.9	8.9	3.0	8.3	2.9
Logical Memory Delayed <sup>7</sup>						
(mean, SD)	6.3	3.2	6.6	3.4	6.1	3.0

<sup>1</sup> Range is 0 to 42; higher score indicates greater literacy.

<sup>2</sup> Range is 0 to 30; higher score indicates better global cognition.

<sup>3</sup> Range is 0 to 140; higher score indicates better processing speed.

<sup>4</sup> Range is 0 to 150 seconds; lower score indicates better processing speed.

<sup>5</sup> Range is 0 to 300 seconds; lower score indicates better processing speed.

<sup>6</sup> Range is 0 to 25; higher score indicates better immediate memory.

<sup>7</sup> Range is 0 to 25; higher score indicates better episodic memory.