

Supplementary table 2

Likelihood and primary site of infection, microbial pathogens and findings at onset of sepsis.

Sepsis (n=158)	
<i>Likelihood of infection according to ISF (n %)</i>	
Confirmed infection	44 (27.8 %)
Probable infection	83 (52.5 %)
Possible infection	17 (10.8 %)
Unknown (ISF N/A)	14 (8.9 %)
<i>Primary site of infection, n (%)</i>	
Pneumonia	113 (71.5 %)
Skin and soft tissue	17 (10.8 %)
Unknown (ISF N/A)	14 (8.9 %)
Secondary peritonitis	11 (7.0 %)
Catheter related	2 (1.2 %)
Urinary tract	1 (0.6 %)
<i>Microbial pathogens (n %)</i>	
None confirmed	100 (63.3 %)
Staphylococcus Aureus	15 (9.5 %)
Escherichia Coli	8 (5.1 %)
Haemophilus Influenzae	7 (4.4 %)
Klebsiella Pneumoniae	4 (2.5 %)
Streptococcus Pneumoniae	3 (1.9 %)
Enterobacter Cloacae	3 (1.9 %)
Enterobacter Aerogenes	3 (1.9 %)
Klebsiella Oxytoca	3 (1.9 %)
Proteus Mirabilis	2 (1.3 %)
Acinetobacter Baumannii	2 (1.3 %)
Proteus Mirabilis	2 (1.3 %)
Other	6 (3.8 %)
<i>Microbial finding (n %)</i>	
None confirmed	100 (60.8 %)
PBS or BAL	40 (28.4 %)
Sputum	9 (5.7 %)
Wound, tissue, abscess	6 (3.4 %)
Blood	2 (1.4 %)
Urine	1 (0.7 %)

Supplementary table 2. Likelihood of infection according to ISF classification, primary site of infection according to ISF classification, microbial pathogens, and microbial findings at debute of sepsis. *ISF*, International Sepsis Forum. *PBS*, Protected Brush Specimen. *BAL*, Bronchoalveolar Lavage.