

Genome-wide association studies for production, respiratory disease, and immune-related traits in Landrace pigs

Yoshinobu Uemoto^{1†*}, Kasumi Ichinoseki^{1†}, Toshimi Matsumoto², Nozomi Oka³, Hironori Takamori³, Hiroshi Kadowaki³, Chihiro Kojima-Shibata³, Eisaku Suzuki³, Toshihiro Okamura⁴, Hisashi Aso¹, Haruki Kitazawa¹, Masahiro Satoh¹, Hirohide Uenishi², Keiichi Suzuki¹

¹Graduate School of Agricultural Science, Tohoku University, Sendai, Miyagi 980-8572, Japan.

²Animal Bioregulation Unit, Division of Animal Sciences, Institute of Agrobiological Sciences, National Agriculture and Food Research Organization (NARO), Tsukuba, Ibaraki 305-8634, Japan

³Miyagi Prefecture Animal Industry Experiment Station, Osaki, Miyagi 989-6445, Japan

⁴Institute of Livestock and Grassland Science, NARO, Tsukuba, Ibaraki 305-0901, Japan

†These authors have contributed equally to this work and share first authorship.

*Correspondence author

Supplementary Figure

Figure S1. Average linkage disequilibrium coefficient (r^2) values plotted against intermarker distance for all autosomal chromosomes in Landrace pigs.




Figure S1. Average linkage disequilibrium coefficient (r^2) values plotted against intermarker distance for all autosomal chromosomes in Landrace pigs. The x-axis indicates the distance between single nucleotide polymorphisms (SNPs) and the y-axis indicates the r^2 values between SNPs.