

**Supplemental Table 2: Comparisons of survey responses according to current position**

Variable	Current position		P-value
	Associate consultant or Consultant (N=81)	Senior Associate Consultant (N=22)	
I am open to using telemedicine routinely in the future for consultations			0.50
Strongly disagree	9 (11.1%)	1 (4.5%)	
Somewhat disagree	12 (14.8%)	5 (22.7%)	
Neither agree not disagree	4 (4.9%)	2 (9.1%)	
Somewhat agree	20 (24.7%)	7 (31.8%)	
Strongly agree	36 (44.4%)	7 (31.8%)	
I am open to using telemedicine routinely in the future for return visits			0.90
Strongly disagree	2 (2.5%)	0 (0.0%)	
Somewhat disagree	4 (4.9%)	1 (4.5%)	
Neither agree not disagree	4 (4.9%)	1 (4.5%)	
Somewhat agree	16 (19.8%)	5 (22.7%)	
Strongly agree	55 (67.9%)	15 (68.2%)	
Assuming both are equally appropriate, I prefer to use telemedicine over face-to-face visits			0.82
Strongly disagree	17 (21.3%)	3 (13.6%)	
Somewhat disagree	11 (13.8%)	5 (22.7%)	
Neither agree not disagree	19 (23.8%)	4 (18.2%)	
Somewhat agree	20 (25.0%)	7 (31.8%)	
Strongly agree	13 (16.3%)	3 (13.6%)	
What role has telemedicine played in your experience of burnout			0.61
Greatly contributed to burnout	2 (2.6%)	0 (0.0%)	
Contributed to burnout	7 (9.2%)	2 (11.1%)	
Neither contributed to or improved burnout	43 (56.6%)	12 (66.7%)	
Alleviated burnout symptoms	20 (26.3%)	3 (16.7%)	
Greatly alleviated burnout symptoms	4 (5.3%)	1 (5.6%)	
Overall, my work-life balance has been improved with telemedicine			0.62
Strongly disagree	9 (11.1%)	1 (4.8%)	
Somewhat disagree	10 (12.3%)	2 (9.5%)	
Neither agree nor disagree	32 (39.5%)	11 (52.4%)	
Somewhat agree	21 (25.9%)	3 (14.3%)	
Strongly agree	9 (11.1%)	4 (19.0%)	

P-values result from a Wilcoxon rank sum test.

**Supplemental Table 3: Comparisons of survey responses according to years in practice following residency**

Variable	Years in practice following residency			P-value
	≤10 years (N=41)	11-20 years (N=23)	>20 years (N=39)	
I am open to using telemedicine routinely in the future for consultations				0.28
Strongly disagree	3 (7.3%)	2 (8.7%)	5 (12.8%)	
Somewhat disagree	8 (19.5%)	3 (13.0%)	6 (15.4%)	
Neither agree not disagree	2 (4.9%)	1 (4.3%)	3 (7.7%)	
Somewhat agree	10 (24.4%)	5 (21.7%)	12 (30.8%)	
Strongly agree	18 (43.9%)	12 (52.2%)	13 (33.3%)	
I am open to using telemedicine routinely in the future for return visits				0.19
Strongly disagree	0 (0.0%)	1 (4.3%)	1 (2.6%)	
Somewhat disagree	2 (4.9%)	0 (0.0%)	3 (7.7%)	
Neither agree not disagree	2 (4.9%)	1 (4.3%)	2 (5.1%)	
Somewhat agree	6 (14.6%)	7 (30.4%)	8 (20.5%)	
Strongly agree	31 (75.6%)	14 (60.9%)	25 (64.1%)	
Assuming both are equally appropriate, I prefer to use telemedicine over face-to-face visits				0.040
Strongly disagree	3 (7.5%)	4 (17.4%)	13 (33.3%)	
Somewhat disagree	8 (20.0%)	3 (13.0%)	5 (12.8%)	
Neither agree not disagree	11 (27.5%)	4 (17.4%)	8 (20.5%)	
Somewhat agree	12 (30.0%)	5 (21.7%)	10 (25.6%)	
Strongly agree	6 (15.0%)	7 (30.4%)	3 (7.7%)	
What role has telemedicine played in your experience of burnout				0.69
Greatly contributed to burnout	0 (0.0%)	0 (0.0%)	2 (5.6%)	
Contributed to burnout	6 (15.8%)	2 (10.0%)	1 (2.8%)	
Neither contributed to or improved burnout	23 (60.5%)	10 (50.0%)	22 (61.1%)	
Alleviated burnout symptoms	7 (18.4%)	8 (40.0%)	8 (22.2%)	
Greatly alleviated burnout symptoms	2 (5.3%)	0 (0.0%)	3 (8.3%)	
Overall, my work-life balance has been improved with telemedicine				0.38
Strongly disagree	2 (4.9%)	1 (4.5%)	7 (17.9%)	
Somewhat disagree	5 (12.2%)	3 (13.6%)	4 (10.3%)	
Neither agree nor disagree	21 (51.2%)	8 (36.4%)	14 (35.9%)	
Somewhat agree	9 (22.0%)	6 (27.3%)	9 (23.1%)	
Strongly agree	4 (9.8%)	4 (18.2%)	5 (12.8%)	

P-values result from Spearman's test of correlation. Years of practice following residency was displayed as a 3-level variable for ease of presentation, but was analyzed as a 5-level ordinal variable in all statistical analysis.

**Supplemental Table 4: Comparisons of survey responses according to age**

Variable	Age 31-40 (N=32)	Age 41-50 (N=30)	Age > 50 (N=41)	P-value
I am open to using telemedicine routinely in the future for consultations				0.56
Strongly disagree	3 (9.4%)	2 (6.7%)	5 (12.2%)	
Somewhat disagree	5 (15.6%)	6 (20.0%)	6 (14.6%)	
Neither agree not disagree	2 (6.3%)	2 (6.7%)	2 (4.9%)	
Somewhat agree	7 (21.9%)	8 (26.7%)	12 (29.3%)	
Strongly agree	15 (46.9%)	12 (40.0%)	16 (39.0%)	
I am open to using telemedicine routinely in the future for return visits				0.39
Strongly disagree	0 (0.0%)	0 (0.0%)	2 (4.9%)	
Somewhat disagree	1 (3.1%)	2 (6.7%)	2 (4.9%)	
Neither agree not disagree	1 (3.1%)	2 (6.7%)	2 (4.9%)	
Somewhat agree	6 (18.8%)	7 (23.3%)	8 (19.5%)	
Strongly agree	24 (75.0%)	19 (63.3%)	27 (65.9%)	
Assuming both are equally appropriate, I prefer to use telemedicine over face-to-face visits				0.048
Strongly disagree	2 (6.3%)	5 (17.2%)	13 (31.7%)	
Somewhat disagree	6 (18.8%)	5 (17.2%)	5 (12.2%)	
Neither agree not disagree	10 (31.3%)	5 (17.2%)	8 (19.5%)	
Somewhat agree	8 (25.0%)	8 (27.6%)	11 (26.8%)	
Strongly agree	6 (18.8%)	6 (20.7%)	4 (9.8%)	
What role has telemedicine played in your experience of burnout				0.51
Greatly contributed to burnout	0 (0.0%)	0 (0.0%)	2 (5.3%)	
Contributed to burnout	5 (16.7%)	2 (7.7%)	2 (5.3%)	
Neither contributed to or improved burnout	17 (56.7%)	17 (65.4%)	21 (55.3%)	
Alleviated burnout symptoms	6 (20.0%)	7 (26.9%)	10 (26.3%)	
Greatly alleviated burnout symptoms	2 (6.7%)	0 (0.0%)	3 (7.9%)	
Overall, my work-life balance has been improved with telemedicine				0.28
Strongly disagree	1 (3.1%)	1 (3.4%)	8 (19.5%)	
Somewhat disagree	6 (18.8%)	2 (6.9%)	4 (9.8%)	
Neither agree nor disagree	14 (43.8%)	14 (48.3%)	15 (36.6%)	
Somewhat agree	7 (21.9%)	8 (27.6%)	9 (22.0%)	
Strongly agree	4 (12.5%)	4 (13.8%)	5 (12.2%)	

P-values result from Spearman's test of correlation. Age was displayed as a 3-level variable for ease of presentation, but was analyzed as a 4-level ordinal variable in all statistical analysis.

**Supplemental Table 5: Comparisons of survey responses according to specialty**

Variable	Specialty				P-value
	Internal/Family Medicine (N=23)	Internal Medicinal Subspecialty (N=42)	Surgery (N=18)	Other (N=20)	
I am open to using telemedicine routinely in the future for consultations					0.14
Strongly disagree	1 (4.3%)	2 (4.8%)	1 (5.6%)	6 (30.0%)	
Somewhat disagree	0 (0.0%)	8 (19.0%)	7 (38.9%)	2 (10.0%)	
Neither agree not disagree	3 (13.0%)	2 (4.8%)	0 (0.0%)	1 (5.0%)	
Somewhat agree	7 (30.4%)	13 (31.0%)	2 (11.1%)	5 (25.0%)	
Strongly agree	12 (52.2%)	17 (40.5%)	8 (44.4%)	6 (30.0%)	
I am open to using telemedicine routinely in the future for return visits					0.058
Strongly disagree	0 (0.0%)	1 (2.4%)	1 (5.6%)	0 (0.0%)	
Somewhat disagree	0 (0.0%)	2 (4.8%)	1 (5.6%)	2 (10.0%)	
Neither agree not disagree	0 (0.0%)	3 (7.1%)	2 (11.1%)	0 (0.0%)	
Somewhat agree	3 (13.0%)	9 (21.4%)	5 (27.8%)	4 (20.0%)	
Strongly agree	20 (87.0%)	27 (64.3%)	9 (50.0%)	14 (70.0%)	
Assuming both are equally appropriate, I prefer to use telemedicine over face-to-face visits					0.001
Strongly disagree	0 (0.0%)	8 (19.5%)	7 (38.9%)	5 (25.0%)	
Somewhat disagree	2 (8.7%)	6 (14.6%)	5 (27.8%)	3 (15.0%)	
Neither agree not disagree	5 (21.7%)	11 (26.8%)	1 (5.6%)	6 (30.0%)	
Somewhat agree	8 (34.8%)	10 (24.4%)	4 (22.2%)	5 (25.0%)	
Strongly agree	8 (34.8%)	6 (14.6%)	1 (5.6%)	1 (5.0%)	
What role has telemedicine played in your experience of burnout					0.006
Greatly contributed to burnout	0 (0.0%)	0 (0.0%)	1 (6.3%)	1 (5.3%)	
Contributed to burnout	0 (0.0%)	4 (10.8%)	4 (25.0%)	1 (5.3%)	
Neither contributed to or improved burnout	11 (50.0%)	23 (62.2%)	10 (62.5%)	11 (57.9%)	
Alleviated burnout symptoms	9 (40.9%)	9 (24.3%)	0 (0.0%)	5 (26.3%)	
Greatly alleviated burnout symptoms	2 (9.1%)	1 (2.7%)	1 (6.3%)	1 (5.3%)	

Overall, my work-life balance has been improved with telemedicine

<0.001

Strongly disagree	0 (0.0%)	2 (4.9%)	4 (22.2%)	4 (20.0%)
Somewhat disagree	1 (4.3%)	4 (9.8%)	4 (22.2%)	3 (15.0%)
Neither agree nor disagree	7 (30.4%)	22 (53.7%)	9 (50.0%)	5 (25.0%)
Somewhat agree	9 (39.1%)	9 (22.0%)	0 (0.0%)	6 (30.0%)
Strongly agree	6 (26.1%)	4 (9.8%)	1 (5.6%)	2 (10.0%)

---

P-values result from a Kruskal-Wallis rank sum test.

---