

Supplemental Online Content

Izadi Z, Li J, Evans M, et al. Socioeconomic disparities in functional status in a national sample of patients with rheumatoid arthritis. *JAMA Netw Open*. 2021;4(8):e2119400. doi:10.1001/jamanetworkopen.2021.19400

eTable. Comparison of Patient Demographic Characteristics Across the Area Deprivation Index Quintiles

This supplemental material has been provided by the authors to give readers additional information about their work.

eTable. Comparison of Patient Demographic Characteristics Across the Area Deprivation Index Quintiles						
Cross-sectional Cohort: Patients with RA and at least 1 FS scores documented during the study period, N=83,965.						
	1st ADI Quintile Highest SES level	2nd ADI Quintile	3rd ADI Quintile	4th ADI Quintile	5th ADI Quintile Lowest SES level	P*
Age, mean (SD)	63.87 (14.1)	63.33 (14.0)	63.32 (13.7)	63.18 (13.5)	63.32 (13.2)	<0.05
Female	12853 (76.9%)	12779 (77.8%)	12895 (76.6%)	13282 (76.6%)	12840 (77.2%)	<0.05
Race/ethnicity						<0.05
Non-Hispanic White	12157 (72.7%)	12133 (73.9%)	12478 (74.1%)	12718 (73.3%)	10551 (63.5%)	
Hispanic	715 (4.3%)	888 (5.4%)	771 (4.6%)	803 (4.6%)	1127 (6.8%)	
African American	775 (4.6%)	966 (5.9%)	1132 (6.7%)	1417 (8.2%)	2610 (15.7%)	
Asian	555 (3.3%)	230 (1.4%)	148 (0.9%)	94 (0.5%)	79 (0.5%)	
Other/Mixed	1161 (6.9%)	979 (6.0%)	981 (5.8%)	939 (5.4%)	936 (5.6%)	
Unknown	1355 (8.1%)	1228 (7.5%)	1335 (7.9%)	1380 (8.0%)	1324 (8.0%)	
Longitudinal Cohort: Patients with RA and ≥ 2 FS scores documented during the study period at least 12-months apart, N=35,385.						
	1st ADI Quintile Highest SES level	2nd ADI Quintile	3rd ADI Quintile	4th ADI Quintile	5th ADI Quintile Lowest SES level	P*
Age, mean (SD)	64.00 (13.5)	63.79 (13.5)	63.60 (13.0)	63.56 (12.9)	63.77 (12.6)	0.29
Female	5584 (77.8%)	5650 (78.9%)	5375 (76.4%)	5346 (76.7%)	5439 (77.2%)	<0.05
Race/ethnicity						<0.05
Non-Hispanic White	5506 (76.7%)	5433 (75.9%)	5300 (75.4%)	5209 (74.7%)	4672 (66.3%)	
Hispanic	284 (4.0%)	322 (4.5%)	251 (3.6%)	218 (3.1%)	369 (5.2%)	
African American	388 (5.4%)	435 (6.1%)	458 (6.5%)	564 (8.1%)	1031 (14.6%)	
Asian	250 (3.5%)	109 (1.5%)	70 (1.0%)	38 (0.6%)	28 (0.4%)	
Other/Mixed	385 (5.4%)	374 (5.2%)	367 (5.2%)	333 (4.8%)	343 (4.9%)	
Unknown	362 (5.1%)	487 (6.8%)	588 (8.4%)	609 (8.7%)	602 (8.6%)	
*Unadjusted p-values were obtained using Chi-squared test for categorical variables and one-way ANOVA test for continuous variables. RA: Rheumatoid arthritis; FS: Functional status;						