

Supplemental Table B: Means and Ranges of the Baseline Measures

Construct/Measure	Mean or Prevalence Rate	Overall Range (Sample Range, if applicable)
Age	20.47 years	18-25 years
Female Sex	65.1% female	not applicable
Race/Ethnicity	53.5% white, non-Hispanic 30.2% white, Hispanic 7.0% black, non-Hispanic 4.7% Asian, non-Hispanic 2.3% American Indian, non-Hispanic 2.5% multiracial	not applicable
Grade Point Average	3.14	0-4 (2.39-4.00)
ADHD Symptom Level <sup>a</sup>	61.51	18-90 (39-78)
Negative Urgency <sup>b</sup>	9.84	4-16 (5-15)
Positive Urgency <sup>b</sup>	11.40	4-16 (4-16)
Lack of Perseverance <sup>b</sup>	7.47	4-16 (4-13)
Sensation Seeking <sup>b</sup>	8.56	4-16 (4-16)
Lack of Premeditation <sup>b</sup>	7.74	4-16 (4-14)
Delay Discounting <sup>c</sup>	0.030	0-1 (0.00016-0.249)
Lifetime Alcohol and PSM Co-ingestion <sup>d</sup>	76.7%	not applicable
Lifetime Non-oral PSM <sup>d</sup>	46.5%	not applicable
Lifetime Recreational PSM Motives <sup>d</sup>	41.9%	not applicable

PSM = Prescription Stimulant Misuse

<sup>a</sup> via the Adult ADHD Self-Report Scale (ASRS) screener

<sup>b</sup> via the Short UPPS-P Impulsive Behavior scale (SUPPS-P)

<sup>c</sup> via the 27-item Monetary Choice Questionnaire from Kirby et al., 1999

<sup>d</sup> via the Stimulant Survey Questionnaire (SSQ)