

SUPPLEMENTAL MATERIAL

Supplemental Table I. Predictors of VOM non-identifiability

Parameter	Univariable model		Multivariable model	
	OR (95% CI)	p-value	OR (95% CI)	p-value
Age	1.01 (0.98-1.05)	0.46		
Male	1.62 (0.78-3.80)	0.21		
Body mass index	1.07 (1.01-1.13)	0.026	1.07 (1.00-1.14)	0.058
Heart failure	0.24 (0.039-0.79)	0.015	0.31 (0.050-1.06)	0.064
Hypertension	0.95 (0.52-1.72)	0.87		
Diabetes mellitus	1.75 (0.77-3.60)	0.17		
Stroke	0.88 (0.26-2.28)	0.82		
Structural heart disease	1.15 (0.62-2.08)	0.66		
Cardiac resynchronization therapy	0.94 (0.051-4.79)	0.95		
Previous open chest surgery	0.97 (0.28-2.50)	0.95		
Previous mitral valve surgery	2.19 (0.63-5.92)	0.20		
Previous endocardial MI ablation	3.52 (1.87-6.58)	0.0001	1.29 (0.51-3.11)	0.59
Previous CS ablation	5.45 (2.83-10.39)	<0.0001	4.08 (1.65-10.19)	0.0024

CI: Confidence interval

CS: Coronary sinus

MI: mitral isthmus

OR: Odds ratio

Supplemental Table II. Clinical characteristics of patients with and without cardiac tamponade

	Total (n=634)	Tamponade (+) (n=7)	Tamponade (-) (n=627)	p-value
Age, years	63.4±10.2	70.4±4.6	63.3±10.2	0.0059
Male, n (%)	476 (75.1)	4 (57.1)	472 (75.3)	0.37
Body mass index, kg/m ²	28.0±4.7	28.9±7.8	28.0±4.6	0.77
Heart failure, n (%)	98 (15.5)	2 (28.6)	96 (15.3)	0.30
Hypertension, n (%)	306 (48.3)	3 (42.9)	303 (48.3)	1.0
Diabetes mellitus, n (%)	76 (12.0)	2 (28.6)	74 (11.8)	0.20
Stroke, n (%)	62 (9.8)	0 (0)	62 (9.9)	1.0
Structural heart disease, n (%)	246 (38.8)	3 (42.9)	243 (38.8)	1.0
History of open chest surgery, n (%)	55 (8.7)	1 (14.3)	54 (8.6)	0.47
History of mitral valve surgery, n (%)	27 (4.3)	1 (14.3)	26 (4.2)	0.26
VOM staining, n (%)	507 (80.0)	6 (85.7)	501 (79.9)	1.0
VOM dissection, n (%)	68 (10.7)	1 (14.3)	67 (10.7)	0.55
VOM perforation, n (%)	14 (2.2)	2 (28.6)	12 (1.9)	0.0089

VOM: Vein of Marshall