Reference	Methods used for test performance estimation	Location	Species	RBT: Se	RBT: Sp	cELISA: Se	cELISA: Sp	Sample size
Marín <i>et al</i> ., 1999 [1]	Se: Culture positive animals (naturally infected)	Unknown	S	1	1	-	-	115
Minas <i>et al</i> ., 2005 [2]	Se: Culture positive animals (naturally infected)	Greece	S	0.976	0.981	-	-	1017
Burriel <i>et al.</i> , 2004 [3]	Se: Vaccinated animals	Greece	SG	-	-	0.885	0.985	383
Minas <i>et al</i> ., 2007 [4]	Se: Serology positive animals from culture confirmed flocks	Greece	SG	0.866	0.976	-	-	4253
Ramírez-Pfeiffer <i>et al</i> ., 2008 [5]	Se: Animals from area with flock abortions and human cases/PCR or serology positive animals from culture confirmed flocks	Mexico & Canada	G	0.997	0.327	-	-	534
McGiven <i>et al.</i> , 2009 [6]	Se: Culture positive animals (naturally or experimentally infected)/Serology positive animals from culture confirmed flock/Animals from suspected outbreak flock in endemic area	UK & Unknown	SG	-	-	1	0.996	281
Thompson <i>et al.</i> , 2009 [7]	Se: Culture positive animals (naturally or experimentally infected)/Serology positive animals from culture confirmed flock/Animals from suspected outbreak flock in endemic area	UK & Unknown	SG	-	-	1	1	201
Perrett <i>et al</i> ., 2010 [8]	Se: Experimentally infected animals	UK	S G	-	-	1 1	0.998 1	484 2023
Weighted mean				0.899	0.919	0.987	0.998	

S3 Table. Mean sensitivity and specificity estimates for the RBT and cELISA in sheep and goats taken from a literature search of global scientific publications from 1999-2019.

Se is sensitivity. Specificity was estimated in each study using samples collected from *Brucella* free animals. S is sheep, G is goats and SG is sheep and goats. RBT is the modified Rose Bengal plate test using the 3:1 serum to antigen ratio. cELISA is the Animal and Plant Health Agency, UK, competitive enzyme-linked immunosorbent assay. Weighted mean is the mean value weighted by sample size.

References

- Marín CM, Moreno E, Moriyón I, Díaz R, Blasco JM. Performance of competitive and indirect enzyme-linked immunosorbent assays, gel immunoprecipitation with native hapten polysaccharide, and standard serological tests in diagnosis of sheep brucellosis. Clin Diagn Lab Immunol. 1999;6: 269–272. doi: 10.1128/CDLI.6.2.269-272.1999.
- Minas A, Stournara A, Minas M, Papaioannou A, Krikelis V, Tselepidis S. Validation of fluorescence polarization assay (FPA) and comparison with other tests used for diagnosis of *B. melitensis* infection in sheep. Vet Microbiol. 2005;111: 211–221. doi: 10.1016/j.vetmic.2005.10.009.
- 3. Burriel AR, Christodoulopoulos G, Bisias G, Fthenakis GC. Comparison of fluorescence polarization assay, indirect ELISA and competitive ELISA methods for diagnosis of *Brucella melitensis*-infection in small ruminants. Small Rumin Res. 2004;54: 243–247. doi: 10.1016/j.smallrumres.2003.12.001.
- Minas A, Stournara A, Minas M, Stack J, Petridou E, Christodoulopoulos G, et al. Validation of a fluorescence polarization assay (FPA) performed in microplates and comparison with other tests used for diagnosing B. melitensis infection in sheep and goats. J Immunol Methods. 2007;320: 94–103. doi: 10.1016/j.jim.2006.12.008.
- Ramírez-Pfeiffer C, Díaz-Aparicio E, Gomez-Flores R, Rodríguez-Padilla C, Morales-Loredo A, Álvarez-Ojeda G. Use of the *Brucella melitensis* Native Hapten To Diagnose Brucellosis in Goats by a Rapid, Simple, and Specific Fluorescence Polarization Assay. Clin Vaccine Immunol. 2008;15: 911–915. doi: 10.1128/CVI.00046-08.
- 6. McGiven JA, Thompson IJ, Commander NJ, Stack JA. Time-Resolved Fluorescent Resonance Energy Transfer Assay for Simple and Rapid Detection of Anti- Brucella Antibodies in Ruminant Serum Samples. J Clin Microbiol. 2009;47: 3098–3107. doi: 10.1128/JCM.00919-09.
- 7. Thompson I, Mcgiven J, Sawyer J, Thirlwall R, Commander N, Stack J. Competitive Electrochemiluminescence Wash and No-Wash Immunoassays for Detection of Serum Antibodies to Smooth *Brucella* Strains. Clin Vaccine Immunol. 2009;16: 765–771. doi: 10.1128/CVI.00006-09.
- 8. Perrett LL, McGiven JA, Brew SD, Stack JA. Evaluation of competitive ELISA for detection of antibodies to *Brucella* infection in domestic animals. Croat Med J. 2010;51: 314–319. doi: 10.3325/cmj.2010.51.314.