Supplementary file 11. Forest plot depicting the effect of combat exposure with PTSD, stratified for study design.

				Odds Ratio	Odds Ratio
Study or Subgroup	log[Odds Ratio]	SE	Weight	IV, Random, 95% CI	IV, Random, 95% CI
3.2.1 Retrospective design					
Connorton, 2011	2.1	0.35	7.1%	8.17 [4.11, 16.22]	_
Green, 2016	0.03	0.02	14.0%	1.03 [0.99, 1.07]	+
Stevelink, 2018	0.7	0.23	9.9%	2.01 [1.28, 3.16]	
Wittchen, 2012	1.89	0.95	1.7%	6.62 [1.03, 42.60]	
Subtotal (95% CI)			32.8%	2.82 [1.08, 7.36]	-
Heterogeneity: Tau ² = 0.79; Chi ² = 46.90, df = 3 (P < 0.00001); I ² = 94%					
Test for overall effect	: Z = 2.11 (P = 0.03)	I			
3.2.2 Prospective de	sign				
Cukor, 2011	0.27	0.07	13.6%	1.31 [1.14, 1.50]	-
Goodwin, 2012	0.69	0.4	6.2%	1.99 [0.91, 4.37]	
Harvey, 2012	0.98	0.29	8.4%	2.66 [1.51, 4.70]	_ _
MacGregor, 2015	0.51	0.19	10.9%	1.67 [1.15, 2.42]	
Osorio, 2018	0.3	0.15	11.9%	1.35 [1.01, 1.81]	
Polusny, 2011	0.85	0.26	9.1%	2.34 [1.41, 3.89]	 −•−
Shea, 2013	0.69	0.35	7.1%	1.99 [1.00, 3.96]	
Subtotal (95% CI)			67.2%	1.66 [1.34, 2.05]	◆
Heterogeneity: Tau ² =	= 0.03; Chi ^z = 12.11	, df = 6	6 (P = 0.0)	6); I² = 50%	
Test for overall effect	: Z = 4.70 (P < 0.000	001)			
Total (95% CI)			100.0%	1.89 [1.46, 2.45]	◆
Heterogeneity: Tau ² = 0.12; Chi ² = 88.86, df = 10 (P < 0.00001); I ² = 89%					
Test for overall effect: Z = 4.79 (P < 0.00001)					
Test for subgroup differences: Chi ² = 1.11, df = 1 (P = 0.29), l ² = 10.2%					

1