Supplementary file 14. Forest plot depicting the effect of combat exposure with PTSD, stratified for type of PTSD ascertainment.

Supplementary me 1 m	. or cot prot depretin	.6		Odds Ratio	Odds Ratio
Study or Subgroup	log[Odds Ratio]	SE	Weight	IV, Random, 95% CI	IV, Random, 95% CI
4.2.1 Probable PTSD					
Goodwin, 2012	0.69	0.4	6.2%	1.99 [0.91, 4.37]	<del>  • </del>
Harvey, 2012	0.98	0.29	8.4%	2.66 [1.51, 4.70]	<del></del>
Osorio, 2018	0.3	0.15	11.9%	1.35 [1.01, 1.81]	<del> </del>
Polusny, 2011	0.85	0.26	9.1%	2.34 [1.41, 3.89]	<del></del>
Stevelink, 2018	0.7	0.23	9.9%	2.01 [1.28, 3.16]	
Subtotal (95% CI)			45.5%	1.90 [1.44, 2.52]	◆
Heterogeneity: Tau² = 0.04; Chi² = 6.81, df = 4 (P = 0.15); l² = 41%					
Test for overall effect:	Z= 4.48 (P < 0.00)	001)			
4.2.2 Diagnosed PTS	D				
Connorton, 2011	2.1	0.35	7.1%	8.17 [4.11, 16.22]	
Cukor, 2011	0.27	0.07	13.6%	1.31 [1.14, 1.50]	
Green, 2016	0.03	0.02	14.0%	1.03 [0.99, 1.07]	<u>†</u>
MacGregor, 2015	0.51	0.19	10.9%	1.67 [1.15, 2.42]	
Shea, 2013	0.69	0.35	7.1%	1.99 [1.00, 3.96]	
Wittchen, 2012	1.89	0.95	1.7%	6.62 [1.03, 42.60]	
Subtotal (95% CI)			54.5%	1.80 [1.28, 2.54]	•
Heterogeneity: Tau² = 0.11; Chi² = 57.78, df = 5 (P < 0.00001); I² = 91%					
Test for overall effect:	Z = 3.39 (P = 0.00)	07)			
Total (95% CI)			100.0%	1.89 [1.46, 2.45]	•
Heterogeneity: Tau² = 0.12; Chi² = 88.86, df = 10 (P < 0.00001); I² = 89%					
Test for overall effect: $Z = 4.79$ (P < 0.00001)					
Test for subgroup differences: Chi <sup>2</sup> = 0.06, df = 1 (P = 0.81), $I^2$ = 0%					