

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Changes in Latina Women's Exposure to Cleaning Chemicals Associated with Switching from Conventional to "Green" Household Cleaning Products: The LUCIR Intervention Study

Kim G. Harley, Lucia Calderon, James E.S. Nolan, Randy Maddalena, Marion Russell, Katherine Roman, Stephanie Mayo-Burgos, Jessica Cabrera, Norma Morga, and Asa Bradman

Table of Contents

Table S1. "Green" products provided to participants by researchers, LUCIR Study, Salinas, California, 2019.

Table S2. Distribution of personal air concentrations of select VOCs ($\mu\text{g}/\text{m}^3$) and SVOCs (ng/m^3) among women using regular cleaning products (pre-intervention visit) and green cleaning products (post-intervention visit), LUCIR Study, Salinas, California, 2019.

Table S3. Most frequently used cleaning products among participants in pre-intervention visit, LUCIR Study, Salinas, California, 2019.