

Additional information file
for
Predictors of help-seeking behaviour in people with mental health problems: a 3-year prospective
community study

**Carolin M. Doll^{1,2}, Chantal Michel³, Marlene Rosen², Naweed Osmann¹, Benno G. Schimmelmann^{3,4},
Frauke Schultze-Lutter^{1,3,5}**

1 Department of Psychiatry and Psychotherapy, Medical Faculty, Heinrich-Heine-University, Düsseldorf, Germany. Electronic address: carolin.doll@uk-koeln.de

2 Department of Psychiatry and Psychotherapy, Faculty of Medicine and University Hospital Cologne, University of Cologne, Cologne, Germany.

3 University Hospital of Child and Adolescent Psychiatry and Psychotherapy, University of Bern, Bern, Switzerland.

4 University Hospital of Child and Adolescent Psychiatry, University Hospital Hamburg-Eppendorf, Hamburg, Germany.

5 Department of Psychology and Mental Health, Faculty of Psychology, Airlangga University, Surabaya, Indonesia

Correspondence author:

Carolin Doll, M.Sc.


Department of Psychiatry and Psychotherapy

Faculty of Heinrich-Heine University and LVR clinic Düsseldorf

Bergische Landstr. 2, 40629 Düsseldorf

Email: carolin.doll@uk-koeln.de, phone: +49221 478 7225

ORCID: <https://orcid.org/0000-0002-4267-1668>



eFig. 2.

Final model of only significant associations between stigma, assumptions about help-seeking and healthcare utilization with standardized path coefficients (N=307).

Model fit indices: $\chi^2_{(27)}=32.174$ with $p=0.226$, CFI=0.985; SRMR=0.047; RMSEA=0.025 (90% CIs=0.000, 0.054).

* $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$; explained variance (R^2) for each endogenous variable in italics. In brackets, Odds Ratios for the endogenous variable “help-seeking within 3 years past baseline” are provided. Manifest variables are represented in rectangles, latent ones in ovals. Solid lines indicate significant paths, dashed lines indicate non-significant paths; in doing so, grey indicates positive, black negative correlations.