

Supplementary material. The raw data of the significant continuous variables

| Reviewed articles | New-onset POAF (n) | Sinus rhythm (n) | Age (mean, SD) | | preOP SCr[mg/dL] (mean, SD) | | Hemoglobin (mean,SD) | | preOP LVEF[%] (mean, SD) | | CBP time[min] (mean, SD) | |
|------------------------------|--------------------|------------------|----------------|---------------|-----------------------------|-------------|----------------------|--------------|--------------------------|---------------|--------------------------|---------------|
| | | | AF | SR | AF | SR | AF | SR | AF | SR | AF | SR |
| Mendes et al. (1995) [21] | 57 | 111 | ☐ | ☐ | ☐ | ☐ | ☐ | ☐ | ☐ | ☐ | ☐ | ☐ |
| Cerillo et al. (2003) [22] | 33 | 74 | 70.2 ± 9.4 | 65.8 ± 9.8 | ☐ | ☐ | ☐ | ☐ | 50.5 ± 9.0 | 53.9 ± 9.7 | 94.4 ± 23.9 | 96.1 ± 29.5 |
| Zangrillo et al. (2004) [23] | 33 | 127 | 68.0 ± 8.8 | 64.0 ± 8.7 | ☐ | ☐ | 14 ± 1.7 | 14 ± 2.6 | ☐ | ☐ | ☐ | ☐ |
| Akazawa et al. (2008) [24] | 26 | 124 | 71.0 ± 8.0 | 66.0 ± 10.0 | ☐ | ☐ | ☐ | ☐ | 49.0 ± 14.0 | 57.0 ± 12.0 | ☐ | ☐ |
| Wang et al. (2012) [25] | 60 | 137 | 71.0 ± 11.0 | 64.0 ± 13.0 | 2.1 ± 2.1 | 1.8 ± 2.0 | ☐ | ☐ | 41.0 ± 11.0 | 48.0 ± 11.0 | ☐ | ☐ |
| Koolen et al. (2013) [26] | 1080 | 2068 | 69.8 ± 8.7 | 64.7 ± 9.8 | 1.12 ± 0.80 | 1.03 ± 0.68 | 8.6 ± 1.0 | 8.7 ± 1.0 | ☐ | ☐ | ☐ | ☐ |
| Tsai et al. (2015) [27] | 126 | 140 | 69.90 ± 11.62 | 61.84 ± 10.76 | ☐ | ☐ | ☐ | ☐ | 48.94 ± 15.19 | 58.19 ± 13.02 | 113.24 ± 58.88 | 98.86 ± 55.01 |
| Vlahou et al. (2016) [28] | 111 | 335 | 68.08 ± 8.5 | 63.05 ± 9.4 | ☐ | ☐ | ☐ | ☐ | 52.8 ± 12.6 | 53.6 ± 10.6 | 97.8 ± 36.1 | 91.1 ± 34.1 |
| Daie et al. (2018) [29] | 29 | 127 | 63.89 ± 8.91 | 61.19 ± 8.90 | 1.00 ± 0.36 | 1.00 ± 0.28 | 13.42 ± 1.35 | 13.68 ± 1.72 | 52.70 ± 10.34 | 54.73 ± 9.32 | ☐ | ☐ |

Abbreviations: POAF; postoperative atrial fibrillation, SD; standard deviation