

## Supplemental Online Content

Falvey JR, Cohen AB, O'Leary JR, Leo-Summers L, Murphy TE, Ferrante LE. Association of social isolation with disability burden and 1-year mortality among older adults with critical illness. *JAMA Intern Med*. Published online September 7, 2021. doi:10.1001/jamainternmed.2021.5022

**eMethods.** Additional Details on Social Isolation Scale Items and Scoring

**eReferences.**

This supplemental material has been provided by the authors to give readers additional information about their work.

## eMethods. Additional Details on Social Isolation Scale Items and Scoring

Below is a description of the survey questions from the National Health and Aging Trends Study (NHATS) used to calculate the social isolation score developed by Pohl and colleagues<sup>1</sup> used in our models. The score ranges from 0 to 6 with higher scores indicating greater isolation.

<b>Social Isolation Domain</b>	<b>NHATS Question</b>	<b>Scoring</b>
Marriage or Partnership	Are you currently married, living with a partner, separated, divorced, widowed, or never married	One point if not married or living with a partner
Connections with family or friends	Looking back over the past year, who are the people you talked with most about often important things?	One point if no family identified One point if no friends identified
	In the last month, did you ever visit with family or friends not living with you, either at home or theirs	One point if answered "no"
Club Participation	In the past month, did you ever participate in clubs, classes, or other organized activities	One point if answered "no"
Religious Service Participation	In the past month, did you ever attend religious services	One point if answered "no"

## eReferences

1. Pohl JS, Cochrane BB, Schepp KG, Woods NF. Measuring social isolation in the national health and aging trends study. *Research in Gerontological Nursing*. 2017;10(6):277-287.