



# Ambient air pollution and low temperature associated with case fatality of COVID-19: A nationwide retrospective cohort study in China


Fei Tian,<sup>1,5</sup> Xiaobo Liu,<sup>2,5</sup> Qingchen Chao,<sup>3</sup> Zhengmin (Min) Qian,<sup>4</sup> Siqi Zhang,<sup>3</sup> Li Qi,<sup>2</sup> Yanlin Niu,<sup>2</sup> Lauren D. Arnold,<sup>4</sup> Shiyu Zhang,<sup>1</sup> Huan Li,<sup>1</sup> Hualiang Lin,<sup>1,\*</sup> and Qiyong Liu<sup>2,\*</sup>

\*Correspondence: [linhualiang@mail.sysu.edu.cn](mailto:linhualiang@mail.sysu.edu.cn) (H.L.); [liuqiyong@icdc.cn](mailto:liuqiyong@icdc.cn) (Q.L.)

Received: December 11, 2020; Accepted: June 16, 2021; Published Online: June 18, 2021; <https://doi.org/10.1016/j.xinn.2021.100139>

© 2021 The Author(s). This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Graphical abstract



## Public summary

- A national retrospective cohort study was performed to quantify the effects of environmental exposure
- Low temperature was observed to be associated with COVID-19 case fatality
- Exposure to ambient air pollution played a nonnegligible part in COVID-19 case fatality
- COVID-19 patients were more susceptible to external environmental stimulus as COVID-19 progressed from the period of symptom onset to diagnosis



# Ambient air pollution and low temperature associated with case fatality of COVID-19: A nationwide retrospective cohort study in China

Fei Tian,<sup>1,5</sup> Xiaobo Liu,<sup>2,5</sup> Qingchen Chao,<sup>3</sup> Zhengmin (Min) Qian,<sup>4</sup> Siqi Zhang,<sup>3</sup> Li Qi,<sup>2</sup> Yanlin Niu,<sup>2</sup> Lauren D. Arnold,<sup>4</sup> Shiyu Zhang,<sup>1</sup> Huan Li,<sup>1</sup> Hualiang Lin,<sup>1,\*</sup> and Qiyong Liu<sup>2,\*</sup>

<sup>1</sup>Department of Epidemiology, School of Public Health, Sun Yat-sen University, Guangzhou 510080, China

<sup>2</sup>State Key Laboratory of Infectious Disease Prevention and Control, Collaborative Innovation Center for Diagnosis and Treatment of Infectious Diseases, WHO Collaborating Centre for Vector Surveillance and Management, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing 102206, China

<sup>3</sup>Laboratory for Climate Studies, National Climate Center, China Meteorological Administration, Beijing 100081, China

<sup>4</sup>Department of Epidemiology and Biostatistics, College for Public Health & Social Justice, Saint Louis University, 3545 Lafayette Avenue, Saint Louis, MO 63104, USA

<sup>5</sup>These authors contributed equally

\*Correspondence: [linhualiang@mail.sysu.edu.cn](mailto:linhualiang@mail.sysu.edu.cn) (H.L.); [liuqiyong@icdc.cn](mailto:liuqiyong@icdc.cn) (Q.L.)

Received: December 11, 2020; Accepted: June 16, 2021; Published Online: June 18, 2021; <https://doi.org/10.1016/j.xinn.2021.100139>

© 2021 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Citation: Tian F., Liu X., Chao Q., et al., (2021). Ambient air pollution and low temperature associated with case fatality of COVID-19: A nationwide retrospective cohort study in China. *The Innovation* 2(3), 100139.

The evidence for the effects of environmental factors on COVID-19 case fatality remains controversial, and it is crucial to understand the role of preventable environmental factors in driving COVID-19 fatality. We thus conducted a nationwide cohort study to estimate the effects of environmental factors (temperature, particulate matter [PM<sub>2.5</sub>, PM<sub>10</sub>], sulfur dioxide [SO<sub>2</sub>], nitrogen dioxide [NO<sub>2</sub>], and ozone [O<sub>3</sub>]) on COVID-19 case fatality. A total of 71,808 confirmed COVID-19 cases were identified and followed up for their vital status through April 25, 2020. Exposures to ambient air pollution and temperature were estimated by linking the city- and county-level monitoring data to the residential community of each participant. For each participant, two windows were defined: the period from symptom onset to diagnosis (exposure window I) and the period from diagnosis date to date of death/recovery or end of the study period (exposure window II). Cox proportional hazards models were used to estimate the associations between these environmental factors and COVID-19 case fatality. COVID-19 case fatality increased in association with environmental factors for the two exposure windows. For example, each 10 µg/m<sup>3</sup> increase in PM<sub>2.5</sub>, PM<sub>10</sub>, O<sub>3</sub>, and NO<sub>2</sub> in window I was associated with a hazard ratio of 1.11 (95% CI 1.09, 1.13), 1.10 (95% CI 1.08, 1.13), 1.09 (95% CI 1.03, 1.14), and 1.27 (95% CI 1.19, 1.35) for COVID-19 fatality, respectively. A significant effect was also observed for low temperature, with a hazard ratio of 1.03 (95% CI 1.01, 1.04) for COVID-19 case fatality per 1°C decrease. Subgroup analysis indicated that these effects were stronger in the elderly, as well as in those with mild symptoms and living in Wuhan or Hubei. Overall, the sensitivity analyses also yielded consistent estimates. Short-term exposure to ambient air pollution and low temperature during the illness would play a nonnegligible part in causing case fatality due to COVID-19. Reduced exposures to high concentrations of PM<sub>2.5</sub>, PM<sub>10</sub>, O<sub>3</sub>, SO<sub>2</sub>, and NO<sub>2</sub> and low temperature would help improve the prognosis and reduce public health burden.

**Keywords:** temperature; air pollution; COVID-19; fatality; cohort study

## INTRODUCTION

At the end of 2019, an outbreak of novel coronavirus (SARS-CoV-2) swept across the globe.<sup>1–3</sup> Although various control measures have been implemented, such as stay-at-home orders, quarantine, social distancing and shielding, the global pandemic continues, and the numbers of cases and deaths still increase. It is therefore critical to identify key modifiable risk factors that affect the case fatality of COVID-19.<sup>4–7</sup>

Environmental factors, especially air pollution and temperature, have been widely identified to be associated with increasing risk of adverse health outcomes, such as for ischemic heart disease, stroke, diabetes, chronic obstructive pulmonary disease, and respiratory infection.<sup>8–11</sup> Emerging evidence indicates potential links between exposure to polluted air and COVID-19 mortality and fatality.<sup>12,13</sup> Recent epidemiological studies indicated that higher concentrations of pollutants were associated with worse outcome of COVID-19. For example, a multicity study from China illustrated that long-term exposure to higher levels of particulate matter was associated with increased COVID-19 case fatality.<sup>14</sup> Wu et al. performed an ecological regression analysis and observed positive effects of higher historical PM<sub>2.5</sub> exposures on county-level COVID-19 mortality rates in the United States.<sup>15</sup> A hierarchical spatial analysis in England reported adverse effects of NO<sub>2</sub> on COVID-19 mortality, but this was not found for PM<sub>2.5</sub>.<sup>16</sup> In addition, several ecological studies have elucidated detrimental effects of NO<sub>2</sub>, with increasing COVID-19 fatality in North America, Europe, and some developed Asian cities.<sup>13,15,17</sup> However, a systematic review of environment and COVID-19 noted some common limitations, such as ecological fallacy due to the use of an aggregated dataset without detailed personal information and adjustment for the individual-level covariates, and these limitations raise concerns about biased or spurious associations.<sup>18</sup>

Toxicological evidence suggests that short-term or long-term environmental exposures could cause pulmonary inflammation and oxidative stress, altering host immune response to viral infections.<sup>19–21</sup> It is thus reasonable to hypothesize that exposure to ambient air pollutants and unfavorable temperature could exacerbate the condition of COVID-19 and increase its case fatality. This study used a national retrospective cohort design to quantify the effects of short-term exposure to ambient low temperature and air pollution (PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>2</sub>, and O<sub>3</sub>) on COVID-19 case fatality in China.

## RESULTS

Of the 657 cities and 2,862 counties/districts in mainland China, 321 cities and 1,676 counties/districts were affected by COVID-19 infection, accounting for about half of the geographic regions in China. A total of 71,808 confirmed COVID-19 cases were included in this nationwide retrospective cohort study. The distribution of the fatality rates across the 321 cities is illustrated in [Figure S1](#). The average age of study participants was 51.43 years, ranging from 1 to 103 years; 49.60% of the participant were females, and 58.29% of participants were home-related workers. Among the included cases, 3,934 died of COVID-19 during the follow-up (through April 25, 2020), resulting in an overall

**Table 1.** Characteristics of study participants by survival status in China

Variable	Deceased (n = 3,934)	Alive (n = 67,874)	p value <sup>a</sup>
<b>Environmental factors</b>			
<b>Exposure window I<sup>a</sup></b>			
Temperature (°C)	5.26 ± 2.82	5.73 ± 4.11	0.0029
Relative humidity (%)	81.43 ± 9.41	79.18 ± 11.36	<0.0001
PM <sub>2.5</sub> (μg/m <sup>3</sup> )	54.01 ± 14.92	51.87 ± 20.25	<0.0001
PM <sub>10</sub> (μg/m <sup>3</sup> )	62.32 ± 16.33	60.81 ± 22.38	<0.0001
O <sub>3</sub> (μg/m <sup>3</sup> )	54.71 ± 7.73	55.84 ± 8.45	<0.0001
SO <sub>2</sub> (μg/m <sup>3</sup> )	7.43 ± 2.01	7.75 ± 3.49	<0.0001
NO <sub>2</sub> (μg/m <sup>3</sup> )	22.78 ± 5.81	20.34 ± 6.38	<0.0001
<b>Exposure window II<sup>a</sup></b>			
Temperature (°C)	7.27 ± 3.04	16.27 ± 2.06	<0.0001
Relative humidity (%)	79.47 ± 6.30	66.44 ± 10.45	<0.0001
PM <sub>2.5</sub> (μg/m <sup>3</sup> )	44.71 ± 15.62	29.62 ± 7.23	<0.0001
PM <sub>10</sub> (μg/m <sup>3</sup> )	53.95 ± 16.76	55.57 ± 11.31	<0.0001
O <sub>3</sub> (μg/m <sup>3</sup> )	55.13 ± 9.31	80.33 ± 9.82	<0.0001
SO <sub>2</sub> (μg/m <sup>3</sup> )	7.41 ± 1.80	9.46 ± 2.63	<0.0001
NO <sub>2</sub> (μg/m <sup>3</sup> )	20.94 ± 6.17	28.98 ± 7.02	<0.0001
Age group (n, %)			<0.0001
<25 years	6 (0.15)	3,493 (5.15)	
25–64 years	1,146 (29.13)	50,111 (73.83)	
≥ 65 years	2,782 (70.72)	14,270 (21.02)	
Sex (n, %)			<0.0001
Male	2,515 (63.93)	33,659 (49.59)	
Female	1,419 (36.07)	34,215 (50.41)	
Days between symptom onset and diagnosis (n, %)			<0.0001
<7 days	1,362 (4.34)	29,995 (95.66)	
7–14 days	1,557 (5.89)	24,859 (94.11)	
≥ 14 days	1,015 (7.23)	13,020 (92.77)	
Occupation (n, %)			<0.0001
Medical-related	21 (0.53)	2,614 (3.85)	
Service-related	15 (0.38)	1,709 (2.52)	
Office worker	226 (5.74)	15,435 (22.74)	
Homemaker	3,226 (82.00)	38,632 (56.92)	
Others	446 (11.34)	9,484 (13.97)	
Residence (n, %)			<0.0001
Permanent	2,476 (62.94)	48,055 (70.80)	
Temporary	1,458 (37.06)	19,819 (29.20)	
Severity (n, %)			<0.0001
Mild	806 (20.49)	28,088 (41.38)	
Moderate	597 (15.18)	28,199 (41.55)	

**Table 1.** Continued

Variable	Deceased (n = 3,934)	Alive (n = 67,874)	p value <sup>b</sup>
Severe	1,285 (32.66)	10,081 (14.85)	
Critical	1,246 (31.67)	1,506 (2.22)	
Hospital transfer (n, %)			0.1754
No	2,927 (74.40)	49,825 (73.41)	
Yes	1,007 (25.60)	18,049 (26.59)	

Note: PM<sub>2.5</sub>, particulate matter with an aerodynamic diameter ≤ 2.5 μm; PM<sub>10</sub>, particulate matter with an aerodynamic diameter ≤ 10 μm; SO<sub>2</sub>, sulfur dioxide; NO<sub>2</sub>, nitrogen dioxide; O<sub>3</sub>, ozone.

<sup>a</sup>Exposure window I represents the mean exposure value from the date of symptom onset to the date of diagnosis; exposure window II represents the mean exposure value from the date of diagnosis to the date of death or the end of the study.

<sup>b</sup>p values were calculated by t test,  $\chi^2$  test or Fisher's exact test, as appropriate.

case fatality rate of 5.48%. There was no loss to follow-up, and the average duration of the follow-up was 81.75 days (SD = 18.01).

Table 1 presents summary statistics for environmental factors using the two exposure windows and individual-level characteristics by survival status. Compared with those who survived, deceased individuals tended to be older (16.31% for age ≥ 65 years versus 0.20% for age <25 years,  $p < 0.001$ ), male (6.95% versus 3.98%,  $p < 0.001$ ), homemakers (7.70% versus 0.80%,  $p < 0.001$ ), and migrants (6.85% versus 4.90%,  $p < 0.001$ ). In critical group that was initially classified, almost half of COVID-19 deaths were identified (45.27%). For exposure window I, there were significant differences in weather conditions between the two groups, with relatively lower temperatures (5.26°C ± 2.82°C) and higher relative humidity (81.43% ± 9.41%) in the deceased group. In addition, there were significant differences in ambient air pollution for the deceased and survived groups, with more highly polluted air for the deceased group compared with the survivor group (54.01 versus 51.87 μg/m<sup>3</sup> for PM<sub>2.5</sub>, 62.32 versus 60.81 μg/m<sup>3</sup> for PM<sub>10</sub>, and 22.78 versus 20.34 μg/m<sup>3</sup> for NO<sub>2</sub>). Significant disparities of environmental factors estimated using window II were also identified. To address concerns about the imbalance of sample size for deceased and survivor groups, we conducted a random resampling of the survivor group to create a subgroup with the same sample size as the deceased group (n = 3,934) and the findings of this resampling analysis were consistent with the full sample results (Table S2). Moreover, we drew maps showing the distribution of air pollution and weather during the relevant period (Figures S2–S7). Table S3 summarizes the Spearman correlation coefficients among key study variables. There are moderate- or low-correlation coefficients between air pollution and meteorological factors (0.01 ≤ |r| ≤ 0.59, except for the high correlation coefficient between PM<sub>2.5</sub> and PM<sub>10</sub>,  $r = 0.96$ ).

Table 2 shows the effect estimates of environmental factors on COVID-19 fatality for two exposure windows (window I and window II). Generally, it was observed that adverse environmental factors were associated with higher fatality rates for COVID-19 in crude and adjusted models. Specifically, the results of the crude model were relatively consistent with those from the adjusted model. We observed significant effects of PM<sub>2.5</sub>, PM<sub>10</sub>, and NO<sub>2</sub>, which were stronger in window II. In the adjusted model for exposure window I, each 1°C decrease in temperature and each 10 μg/m<sup>3</sup> increase in air pollution (PM<sub>2.5</sub>, PM<sub>10</sub>, and NO<sub>2</sub>) concentrations was associated with an HR of 1.03 (95% CI 1.01, 1.04), 1.11 (95% CI 1.09, 1.13), 1.10 (95% CI 1.08, 1.13), and 1.27 (95% CI 1.19, 1.35), respectively, for fatality due to COVID-19.

Figure 1 shows the exposure-response curves of the effects of environmental factors (exposure window I) on COVID-19 fatality. Overall, the temperature curve has a downward trend, although there is a small rise when temperature is below freezing. With the exception of SO<sub>2</sub>, all air pollutants showed curves with upward trends as the concentrations increase; these relationships were monotonic and approximately linear.

The associations of environmental factors with COVID-19 varied by a few characteristics (Table 3). Stronger effects occurred in the elderly, as well as

**Table 2.** Hazard ratios and 95% CIs for case fatalities due to COVID-19 associated with environmental factors of two exposure windows in China

Variable	Hazard ratio (95% CI)				p value <sup>b</sup>
	Exposure window I <sup>a</sup>		Exposure window II <sup>a</sup>		
	Crude <sup>c</sup>	Adjusted <sup>d</sup>	Crude <sup>c</sup>	Adjusted <sup>d</sup>	
Temperature	1.02 (1.01, 1.03)	1.03 (1.01, 1.04)	1.27 (1.26, 1.28)	3.20 (3.15, 3.25)	<0.01
PM <sub>2.5</sub>	1.05 (1.04, 1.07)	1.11 (1.09, 1.13)	1.56 (1.54, 1.58)	1.39 (1.36, 1.42)	<0.01
PM <sub>10</sub>	1.04 (1.02, 1.05)	1.10 (1.08, 1.13)	1.43 (1.42, 1.44)	1.18 (1.16, 1.21)	<0.01
NO <sub>2</sub>	1.69 (1.93, 1.75)	1.27 (1.19, 1.35)	1.22 (1.14, 1.30)	1.37 (1.29, 1.45)	0.08
SO <sub>2</sub>	0.70 (0.62, 0.79)	1.10 (0.95, 1.27)	1.13 (0.91, 1.40)	1.51 (1.22, 1.88)	0.20
O <sub>3</sub>	0.84 (0.81, 0.87)	1.09 (1.03, 1.14)	1.27 (1.21, 1.33)	1.25 (1.20, 1.30)	<0.01

Note: PM<sub>2.5</sub>, particulate matter with an aerodynamic diameter  $\leq 2.5$   $\mu\text{m}$ ; PM<sub>10</sub>, particulate matter with an aerodynamic diameter  $\leq 10$   $\mu\text{m}$ ; SO<sub>2</sub>, sulfur dioxide; NO<sub>2</sub>, nitrogen dioxide; O<sub>3</sub>, ozone.

<sup>a</sup>Exposure window I represents the mean exposure value from the date of symptom onset to the date of diagnosis; exposure window II represents the mean exposure value from the date of diagnosis to the date of death or the end of the study.

<sup>b</sup>Estimated using likelihood ratio test by comparing adjusted HRs for different windows.

<sup>c</sup>Crude model, without any adjustment.

<sup>d</sup>Multivariate model, adjusted for age, sex, occupation, residence, severity of illness, location, transfer history, temporal trend, lockdown, city-level GDP, hospital beds per 1,000 persons, temperature (only for the pollutants), and relative humidity.

those with mild symptoms and living in Wuhan or Hubei. In particular, for an increase of 10  $\mu\text{g}/\text{m}^3$  in air pollutant (PM<sub>2.5</sub>, PM<sub>10</sub>, and NO<sub>2</sub>) concentration, the estimated HRs were slightly larger in those  $\geq 65$  ( $p < 0.01$ ). The HRs for COVID-19 fatality related to each 10  $\mu\text{g}/\text{m}^3$  increase in NO<sub>2</sub> were higher and significant for mild patients compared with severe cases. In addition, almost all significant associations were observed across subgroups of Hubei Province and outside Hubei, although higher effects occurred in Hubei ( $p < 0.05$ ). This relatively consistent pattern was also observed in Wuhan and outside Wuhan. No significant differences were observed in the subgroups before and after city lockdown.

Sensitivity analyses (Table S4) demonstrated that the effects of air pollution remained robust as the degrees of freedom of parameters changed from 2 to 6. For example, the magnitude of HRs for COVID-19 fatality associated with each 10  $\mu\text{g}/\text{m}^3$  increase in PM<sub>10</sub> remained at the level of 1.10. When the association between environmental factors and case fatality due to COVID-19 was explored using the current symptom onset date and diagnosis as well as corresponding lag days, there were significant effects of environmental factors on COVID-19 fatality across lag days (Figures S8 and S9). In addition, the estimated effects of the primary pollutant varied slightly when an adjustment was made for a second pollutant (Table S5). The exception was for NO<sub>2</sub> and O<sub>3</sub>, whose effects were rendered nonsignificant after adjustment for PM<sub>2.5</sub> and PM<sub>10</sub>, respectively. We also obtained similar results for combined exposure windows compared with exposure window II (Table S6).

## DISCUSSION

In this nationwide retrospective cohort study of COVID-19 patients in China, short-term exposure to ambient environmental factors during the illness course, including low temperature and PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub>, SO<sub>2</sub>, and O<sub>3</sub>, contributed to increasing risk of death. Several exposure windows were used to examine the effects, and the estimates remained robust. To the best of our knowledge, this is the first study to explore the association of exposures to key environmental factors and COVID-19 fatality using nationwide data from low- and middle-income countries with high air pollution levels.


Air pollutants were critical environmental drivers in triggering COVID-19 fatality. These findings are consistent with a previous study that provided evidence for an association between higher air pollution and increased risk of dying from SARS.<sup>22</sup> In addition, emerging research has reported that chronic exposure to polluted air might result in a higher risk of dying from COVID-19.<sup>23</sup> A recent study using county-level data for the United States found that an 8% (95% CI 2%, 15%) increase in the COVID-19 death rate was associated with an increase of 1  $\mu\text{g}/\text{m}^3$  in PM<sub>2.5</sub>.<sup>24</sup> Long-term exposure to NO<sub>2</sub> has also been identified as an important contributor to fatality due to COVID-19.<sup>12</sup> The significant associations between other major ambient pollutants (e.g., SO<sub>2</sub> and O<sub>3</sub> levels) and increased COVID-19 deaths were highlighted in a recent study in England.<sup>17</sup> However, a cross-sectional national study in the United States did not observe significant associations of PM<sub>2.5</sub> and O<sub>3</sub> with COVID-19 fatality.<sup>13</sup> In general, our evidence was derived from a nationwide cohort study with a wider exposure range, providing relatively accurate and convincing estimates.

Further, while numerous studies have focused on the link between COVID-19 transmission and temperature,<sup>25,26</sup> the impact of temperature on COVID-19 outcomes has not been widely studied. We observed that lower temperature was associated with a greater HR of COVID-19 fatality. These findings were consistent with one recent study from Wuhan, China, which reported that higher ambient temperature was related to a lower case fatality rate due to COVID-19.<sup>27</sup> However, the role of ambient temperature in COVID-19 fatality might be rather intricate due to the adverse effects of high and low temperatures on human health.<sup>28–30</sup> Other seasonal data are required to determine the overall comprehensive effects of temperature.

Most COVID-19 cases were admitted in hospitals, particularly, Fangcang shelter hospitals in Wuhan, and these were large and temporary hospitals built by converting public venues.<sup>31</sup> In the Fangcang shelter hospitals or infectious disease zones of designated hospitals, the ventilation systems maintained air exchange from outside to inside, ensuring the plausibility and accuracy of exposure assessment. Although the underlying molecular mechanisms of environmental pollutants and temperature on COVID-19 outcomes remain to be determined, prior studies offered some insight as to potential biological plausibility.<sup>19,32–34</sup> Exposure to air pollutants can induce systemic oxidative stress and enhance inflammation, interacting with SARS-CoV-2 (the virus that causes COVID-19) to damage lymphocytes and impair the immune system.<sup>35</sup> Elevated inflammatory factors and immune dysfunction might trigger the cytokine storm syndrome and develop into acute respiratory distress syndrome, which was associated with COVID-19 disease severity and death.<sup>36</sup> In addition, short-term effects of cold temperatures on health mainly can result in restricted immune function and lead to changes in autonomic nervous system function, inflammatory response, and oxidative stress.<sup>37</sup> In these situations, the symptoms of COVID-19 would progress, and the patient's condition would further deteriorate.

This study adds new evidence on the association between environmental factors and COVID-19 fatality. Comprehensive understanding of the role that external environmental factors play in triggering case fatality due to COVID-19 has valuable implications. Although we do not need another reason to reduce exposure to a poor environment, reporting these associations enriches the long list of the health effects of air pollution. More crucial is that these findings should be expanded to change clinical practice or public health programming, including identification of patients with poor prognosis at early stages, guiding resource allocation, and improvement of prognosis. Findings suggest that COVID-19 patients living in regions with high adverse environmental factors may be more vulnerable to complications from the virus, specifically higher risk for death. These environmental factors are key modifiable variables related to COVID-19 fatality. Hence, all COVID-19 patients should pay attention to self-protection to decrease their exposure to environmental stimuli, such as staying in a warmer environment and avoiding severe air pollution outside.

The results of stratified analyses indicated that those over age 65 were more susceptible to COVID-19 death, which was in line with a previous study.<sup>38</sup> We also observed inconsistent associations across subgroups of Hubei Province and outside Hubei. The reason for this may be due to a



**Figure 1. Exposure-response curves for the effects of environmental factors** The exposure-response curves for exposure to ambient temperature (A),  $PM_{2.5}$  (B),  $PM_{10}$  (C),  $NO_2$  (D),  $SO_2$  (E), and  $O_3$  (F) versus case fatality due to COVID-19 (exposure window I) are shown. The solid and dashed lines represent log (hazard ratio) values and the 95% CIs, respectively. Definition of abbreviations: CI, confidence interval;  $PM_{2.5}$ , particulate matter with an aerodynamic diameter  $\leq 2.5 \mu m$ ;  $PM_{10}$ , particulate matter with an aerodynamic diameter  $\leq 10 \mu m$ ;  $SO_2$ , sulfur dioxide;  $NO_2$ , nitrogen dioxide;  $O_3$ , ozone.

high proportion of severe and critical cases and/or insufficient amounts of medical services in Hubei Province due to it being an epicenter. In addition, we observed a stronger effect of environmental exposures in mild cases compared with severe ones, which could be explained by the scenario that among the severe, there was a portion of COVID-19 intensive care cases (about 19.5% of severe cases) likely admitted to the intensive care unit with lower opportunity to be exposed to external environmental factors.<sup>39</sup> However, although patients were admitted to the wards, they were still affected by the external environment exposures, as there is continuous circulation between indoor and outdoor air.

Two exposure windows were used to examine the association of air pollution and temperature with COVID-19 fatality, and the estimates varied somewhat. We observed effects estimation of window II > window I, suggesting that patients were more susceptible to external environmental stimuli as COVID-19 progressed from the period of symptom onset to diagnosis. An underlying mechanism linking this phenomenon might be that the body system and internal environment experienced distinct changes in innate immune response and inflammation activation as the disease progressed, causing the body to enter a more susceptible and vulnerable state.<sup>40,41</sup> Therefore, the cases during this sensitive period should be given higher priority and more care.

### Strengths and limitations

Our study has several strengths: the large sample size of a stable target population, a longitudinal study with individual-level information, clear temporality between exposure and outcomes, the length of the follow-up from onset through vital outcomes, and adjustment for a wide range of covariates using regression models. It is recognized that the effects of environmental drivers on health outcomes mainly depend on the approaches with which they are evaluated. In this study, several exposure windows were used to evaluate environmental short-term effects, which in turn added to the robustness of the findings.

Our study also has some limitations. As this study used exposure data based on measurements from fixed air monitoring stations as the surrogate for individual exposure, exposure misclassification may be one concern. Nevertheless, this exposure assessment approach has been widely used to develop exposure estimations for large air pollution epidemiological studies, and previous studies reported that applying the monitoring values as personal exposure tended to cause underestimation of effects.<sup>42</sup> The observed effects may still contain residual confounding due to unavailability of other individual behavioral risk factors and unmeasured time-varying confounding factors. Although some relevant individual variables, such as underlying health conditions, BMI, and smoking, were not included in the analysis due to data unavailability, we adjusted for clinical severity as one proxy for the unmeasured clinical information.<sup>43</sup> In addition, our cohort was followed until April 25, 2020. This decision was due to data availability, although that time may not be sufficient to capture all outcomes. Moreover, although we have added one binary variable to the model to control for the effects of city lockdown, it is still insufficient to account for the lockdown effects on the associations due to lack of detailed controlled measures across different cities.

### Conclusion

This study suggests that short-term exposure to air pollution and low temperature could increase the case fatality due to COVID-19 in China. Research on how environmental factors may drive COVID-19 case fatality is crucial to guiding clinical practice and public health programming to curb influences related to the outbreak.

### MATERIAL AND METHODS

#### Study population and design

This retrospective cohort study used the National Notifiable Disease Reporting System of China to enroll all confirmed COVID-19 cases from December 8, 2019, through April 15, 2020. Cases were followed up for vital status until April 25, 2020. Eligible

**Table 3.** Subgroup analysis for the associations of COVID-19 fatality with environmental exposure

Variable	Temperature <sup>a</sup>		PM <sub>2.5</sub> <sup>a</sup>		PM <sub>10</sub> <sup>a</sup>		NO <sub>2</sub> <sup>a</sup>		SO <sub>2</sub> <sup>a</sup>		O <sub>3</sub> <sup>a</sup>		
	HR (95% CI)	p value <sup>b</sup>	HR (95% CI)	p value <sup>b</sup>	HR (95% CI)	p value <sup>b</sup>	HR (95% CI)	p value <sup>b</sup>	HR (95% CI)	p value <sup>b</sup>	HR (95% CI)	p value <sup>b</sup>	
Age		0.02		<0.01		<0.01		<0.01		<0.01		0.11	0.58
<65	1.00 (0.97, 1.02)		1.06 (1.02, 1.10)		1.05 (1.01, 1.09)		1.11 (1.00, 1.24)		0.89 (0.68, 1.17)		1.07 (1.00, 1.17)		
≥ 65	1.04 (1.02, 1.06)		1.14 (1.11, 1.17)		1.13 (1.11, 1.16)		1.34 (1.23, 1.45)		1.16 (0.98, 1.38)		1.10 (1.04, 1.16)		
Sex		0.81		0.64		0.52		0.15		0.50		0.32	
Male	1.07 (0.98, 1.17)		1.04 (0.96, 1.14)		1.07 (1.00, 1.17)		1.22 (1.13, 1.33)		1.03 (0.86, 1.23)		1.07 (1.01, 1.13)		
Female	1.09 (0.97, 1.23)		1.07 (1.00, 1.17)		1.12 (1.00, 1.25)		1.35 (1.21, 1.51)		1.14 (0.90, 1.45)		1.12 (1.04, 1.20)		
Severity		<0.01		0.09		0.10		<0.01		0.08		0.39	
Mild	1.13 (1.08, 1.20)		1.18 (1.11, 1.27)		1.16 (1.09, 1.23)		1.62 (1.36, 1.94)		1.91 (1.05, 3.51)		1.18 (1.04, 1.35)		
Severe	1.01 (1.00, 1.03)		1.11 (1.09, 1.13)		1.10 (1.08, 1.12)		1.25 (1.16, 1.34)		1.10 (0.96, 1.27)		1.11 (1.06, 1.17)		
Location		<0.01		<0.01		<0.01		<0.01		0.86		<0.01	
Wuhan	1.15 (1.12, 1.18)		1.23 (1.19, 1.28)		1.19 (1.15, 1.23)		1.44 (1.33, 1.57)		1.12 (0.69, 1.79)		1.19 (1.11, 1.28)		
Non-Wuhan	1.00 (0.98, 1.02)		1.06 (1.03, 1.09)		1.06 (1.03, 1.09)		1.04 (0.93, 1.16)		1.07 (0.92, 1.25)		1.03 (0.96, 1.09)		
Province		<0.01		0.11		0.03		0.03		<0.01		0.07	
Hubei	1.12 (1.09, 1.15)		1.04 (1.00, 1.05)		1.08 (1.03, 1.13)		1.65 (1.55, 1.76)		1.38 (1.12, 1.71)		1.12 (0.96, 1.32)		
Non-Hubei	1.02 (1.00, 1.05)		1.09 (1.04, 1.15)		1.02 (1.00, 1.04)		1.33 (1.11, 1.59)		0.71 (0.57, 0.89)		0.96 (0.92, 1.00)		
Lockdown <sup>c</sup>		0.29		0.11		0.07		0.16		0.06		<0.01	
After	1.05 (1.02, 1.09)		1.07 (1.02, 1.13)		1.06 (1.02, 1.11)		1.17 (1.04, 1.32)		0.86 (0.51, 1.22)		0.93 (0.53, 1.33)		
Before	1.03 (1.01, 1.04)		1.12 (1.09, 1.14)		1.11 (1.08, 1.14)		1.30 (1.19, 1.42)		1.37 (1.16, 1.62)		1.21 (1.14, 1.29)		

Note: HR, hazard ratio; PM<sub>2.5</sub>, particulate matter with an aerodynamic diameter  $\leq 2.5 \mu\text{m}$ ; PM<sub>10</sub>, particulate matter with an aerodynamic diameter  $\leq 10 \mu\text{m}$ ; SO<sub>2</sub>, sulfur dioxide; NO<sub>2</sub>, nitrogen dioxide; O<sub>3</sub>, ozone.

<sup>a</sup>All the environmental factors were estimated using exposure window I (mean exposure value from the date of symptom onset to the date of diagnosis). The effects of environmental factors were estimated using the adjusted model, adjusted for age, sex, temperature (only for the pollutants), relative humidity, occupation, residence, severity of the illness, location, transfer history, temporal trend, and lockdown.

<sup>b</sup>p values were estimated using likelihood ratio test by comparing different subgroups, and  $p < 0.05$  indicates statistical significance.

<sup>c</sup>Lockdown: on January 23, 2020, the central government of China imposed a lockdown in Wuhan and other cities in Hubei.

participants included those: (1) diagnosed as confirmed COVID-19 cases with a positive viral nucleic acid test result on oropharyngeal swab samples in strict accordance with national criteria and (2) who had complete medical and epidemiological information, including severity of the illness, date of symptom onset, and date of diagnosis. A total of 1,752 cases (2.38%) were excluded from 73,560 original cases due to unavailability of environmental exposure data. The remaining 71,808 COVID-19 cases were included in this analysis. General demographic and illness-related characteristics were obtained from the National Notifiable Disease Reporting System of China, including age, sex, birth date, occupation, residential location, severity of the illness, hospital transfer history, date of symptom onset, date of diagnosis, and date of death (for the deceased).

Approval to conduct this study was obtained from the Biomedical Research Ethics Review Committee of Sun Yat-sen University School of Public Health (no. 2020016).

#### Estimation of environmental exposures

Daily ambient air pollution data (PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub>, SO<sub>2</sub>, and O<sub>3</sub>) during the study period were obtained from China's National Real-Time Publishing Platform for Daily Air Quality, which included 1,256 monitors at the end of 2015, with the number in each city ranging from 1 to 17. There is a series of supporting standards and regulations for air-quality monitoring process and choice of location, which could ensure representativeness of the general background levels of air pollution for the population's daily air pollution exposure.<sup>44</sup> Due to the unavailability of each case's detailed residential address, city-wide average data were used to represent exposure based on the exposure windows. Each participant's exposures were estimated by matching residential city with air pollution concentration in the corresponding city. Considering possible differences and susceptibility of short-term exposure to adverse environmental factors at different stages of COVID-19, we divided the exposures into different periods. Exposure window I represents the period of time from symptom onset to diagnosis, indicating the environmental exposure before hospitalization or quarantining. Exposure window II includes the time from date of diagnosis to date of death or recovery

ery or end of the study period and represents participants' environmental exposures during treatment. In addition, we combined the two exposure windows to create one exposure window for the full course of symptomatic disease (i.e., symptom onset to death or recovery or end of the study). This complete exposure window was calculated for each participant to assess overall effects of the environmental factors. The county-level daily temperature ( $^{\circ}\text{C}$ ) and relative humidity (%) were obtained from the National Meteorological Monitoring Center. Mean temperature was then calculated for each case for each of the aforementioned exposure windows.

#### Statistical analysis

Student's t test was used to examine statistical differences for the continuous variables according to the vital status and expressed as mean  $\pm$  standard deviation (SD). Categorical variables were presented with percentages, and chi-square tests were used to evaluate differences. Spearman's correlation coefficients were estimated among study variables to exclude the potential multicollinearity.

Cox proportional hazards models were applied to estimate the association between COVID-19 fatality and the two exposures of air pollution and temperature. Schoenfeld residuals were performed to evaluate the proportional hazards assumption and no violations were observed ( $p > 0.05$ ). Univariate analysis was conducted, and then multivariate models were generated to control for potential confounding factors. Two criteria were used to select covariates for the final adjusted model: (1) variables known or hypothesized to be risk factors for COVID-19 fatality according to the previous study<sup>38</sup> and (2) variables significantly associated with COVID-19 fatality in the univariate analysis (Table S1). Based on these criteria, the following variables were included in the model: sex (male or female), age (<25 years, 25–65 years, and  $\geq 65$  years), occupation (medical-related, service-related, office worker, homemaker, and others), residence (permanent or temporary), clinically diagnosed severity of the illness (mild, moderate, severe, and critical), location (Wuhan or non-Wuhan), transfer to a higher-level hospital (yes or no), and relative humidity and temperature. Nonparametric smoothing was done by way of a natural smoothing splines function for trend on days

included in the model to account for the natural temporal trend of associations and other potential factors related to COVID-19 fatality. The temporal trend, temperature, and relative humidity were characterized by natural cubic splines for their potentially nonlinear relationship, with 3 degrees of freedom for each based on previous studies.<sup>45,46</sup> In addition, to account for the effects of city lockdown, we determined whether the date of symptoms of onset for each case occurred before or after the city lockdown in each city and added one binary variable to the model. The Chinese government committed to paying for all COVID-19-related hospital bills for every patient. In theory, this would eliminate confounding by some health and medical variables related to access to care and resources. Despite this, we included city-level per-capita GDP and hospital beds per 1,000 persons (each as derived from the China Health Statistical Yearbook 2019) in the model to adjust for potential confounding bias. Details of the overall model are illustrated in the supplemental information (Text S1 and S2). We also plotted the exposure-response relationship curves between environmental factors and COVID-19 fatality with natural cubic spline functions with 3 degrees of freedom in the regression model to examine the nonlinear relationship. Hazard ratios (HRs) and 95% confidence intervals (CIs) were computed for COVID-19 fatality associated with per °C decrease in temperature or per 10 µg/m<sup>3</sup> increase in each air pollutant.

### Subgroup analysis

Stratified analyses by sex (male and female), age group (<65 years and ≥ 65 years), location (Wuhan and non-Wuhan), and the clinically diagnosed severity of illness (mild [including the original mild and moderate] and severe [including the original severe and critical]) were performed to examine potential effect modifiers. In addition, considering the potential disparities in medical resources, treatment supplies, and public health interventions by region and time, study participants were divided into those living outside of Hubei Province and within Hubei, before lockdown and after lockdown. Separate Cox models for each subgroup described above were applied to obtain corresponding HRs for environmental factors. The statistical difference between subgroups was calculated as follows:  $E_1 - E_2 \pm 1.96 * \sqrt{(SE_1)^2 + (SE_2)^2}$ .  $E_1$ ,  $E_2$ ,  $SE_1$ , and  $SE_2$  presented the effect estimates and corresponding SDs in two subgroups.<sup>47</sup> Considering the possible type I errors introduced by multiple runs of the likelihood ratio test, we conducted the Bonferroni correction of likelihood ratio test for testing multiple hypotheses to avoid potential type I errors.<sup>48</sup>

### Sensitivity analysis

Several sensitivity analyses were performed to examine the robustness of our estimates. First, we conducted two-pollutant models with adjustment for a second air pollutant to assess the independent effect of the primary one. Second, different degrees of freedom from 2 to 6 for temperature and relative humidity were specified to examine the impact of alternative degrees of freedom on the estimated effects of environmental factors. Finally, other exposure windows were assigned to the participants, including single-day exposure prior to symptom onset date and diagnosis date (lag<sub>0</sub> to lag<sub>3</sub> for air pollution, lag<sub>0</sub> to lag<sub>10</sub> for temperature). Considering the accumulative effects of environmental factors, the associations between the exposures and COVID-19 were examined using multiday lag structures: a moving average of the current day and the previous 1, 2, and 3 or more days (lag<sub>01</sub>, lag<sub>02</sub>, and lag<sub>03</sub> for air pollution; lag<sub>03</sub>, lag<sub>05</sub>, lag<sub>07</sub>, and lag<sub>010</sub> for temperature).

Statistical analyses were performed by R software. A value of two-tailed  $p < 0.05$  was considered statistically significant.

### REFERENCES

- Wang, Y., Liu, Y., Liu, L., et al. (2020). Clinical outcomes in 55 patients with severe acute respiratory syndrome coronavirus 2 who were asymptomatic at hospital admission in Shenzhen, China. *J. Infect Dis.* **221**, 1770–1774. <https://doi.org/10.1093/infdis/jiaa119>.
- Aslam, A., Singh, J., Robilotti, E., et al. (2020). Severe acute respiratory syndrome coronavirus 2 surveillance and exposure in the perioperative setting with universal testing and personal protective equipment policies. *Clin. Infect Dis.* <https://doi.org/10.1093/cid/ciaa1607>.
- Kar, S., and Leszczynski, J. (2020). From animal to human: interspecies analysis provides a novel way of ascertaining and fighting COVID-19. *Innovation* **1**, 100021. <https://doi.org/10.1016/j.xinn.2020.100021>.
- Zheng, X.-y., Guan, W.-j., and Zhong, N.-s. (2021). Clinical characteristics of COVID-19 in developing countries of western pacific: low case-fatality rate unraveled. *Lancet Reg. Health West. Pac.* **6**, 100073. <https://doi.org/10.1016/j.lanwpc.2020.100073>.
- Dai, C.L., Kornilov, S.A., Roper, R.T., et al. (2021). Characteristics and factors associated with coronavirus disease 2019 infection, hospitalization, and mortality across race and ethnicity. *Clin. Infect Dis.* <https://doi.org/10.1093/cid/ciab154>.
- Huh, K., Kim, Y.-E., Ji, W., et al. (2021). Decrease in hospital admissions for respiratory diseases during the COVID-19 pandemic: a nationwide claims study. *Thorax*. <https://doi.org/10.1136/thoraxjnl-2020-216526>.
- Ra, S.H., Lim, J.S., Kim, G.-u., et al. (2021). Upper respiratory viral load in asymptomatic individuals and mildly symptomatic patients with SARS-CoV-2 infection. *Thorax* **76**, 61–63. <https://doi.org/10.1136/thoraxjnl-2020-215042>.
- Yang, Y., Qi, J., Ruan, Z., et al. (2020). Changes in life expectancy of respiratory diseases from attaining daily PM2.5 standard in China: a nationwide observational study. *Innovation* **1**, 100064. <https://doi.org/10.1016/j.xinn.2020.100064>.
- Yu, W., Guo, Y., Shi, L., et al. (2020). The association between long-term exposure to low-level PM2.5 and mortality in the state of Queensland, Australia: a modelling study with the difference-in-differences approach. *PLoS Med.* **17**, e1003141. <https://doi.org/10.1371/journal.pmed.1003141>.
- Zhao, Q., Li, S., Coelho, M., et al. (2019). The association between heatwaves and risk of hospitalization in Brazil: a nationwide time series study between 2000 and 2015. *PLoS Med.* **16**, e1002753. <https://doi.org/10.1371/journal.pmed.1002753>.
- Ruan, Z., Qi, J., Qian, Z., et al. (2021). Disease burden and attributable risk factors of respiratory infections in China from 1990 to 2019. *Lancet Reg. Health West. Pac.* **11**, 100153. <https://doi.org/10.1016/j.lanwpc.2021.100153>.
- Ogen, Y. (2020). Assessing nitrogen dioxide (NO2) levels as a contributing factor to coronavirus (COVID-19) fatality. *Sci. Total Environ.* **726**, 138605. <https://doi.org/10.1016/j.scitotenv.2020.138605>.
- Liang, D., Shi, L., Zhao, J., et al. (2020). Urban air pollution may enhance COVID-19 case-fatality and mortality rates in the United States. *Innovation* **1**, 100047. <https://doi.org/10.1016/j.xinn.2020.100047>.
- Yao, Y., Pan, J., Wang, W., et al. (2020). Association of particulate matter pollution and case fatality rate of COVID-19 in 49 Chinese cities. *Sci. Total Environ.* **741**, 140396. <https://doi.org/10.1016/j.scitotenv.2020.140396>.
- Wu, X., Nethery, R.C., Sabath, M.B., et al. (2020). Air pollution and COVID-19 mortality in the United States: strengths and limitations of an ecological regression analysis. *Sci. Adv.* **6**. <https://doi.org/10.1126/sciadv.abd4049>.
- Konstantinoudis, G., Padellini, T., Bennett, J., et al. (2021). Long-term exposure to air pollution and COVID-19 mortality in England: a hierarchical spatial analysis. *Environ. Int.* **146**, 106316. <https://doi.org/10.1016/j.envint.2020.106316>.
- Travaglio, M., Yu, Y., Popovic, R., et al. (2021). Links between air pollution and COVID-19 in England. *Environ. Pollut.* **268**, 115859. <https://doi.org/10.1016/j.envpol.2020.115859>.
- Villeneuve, P.J., and Goldberg, M.S. (2020). Methodological considerations for epidemiological studies of air pollution and the SARS and COVID-19 coronavirus outbreaks. *Environ. Health Perspect.* **128**, 95001. <https://doi.org/10.1289/EHP7411>.
- Wu, S., Ni, Y., Li, H., et al. (2016). Short-term exposure to high ambient air pollution increases airway inflammation and respiratory symptoms in chronic obstructive pulmonary disease patients in Beijing, China. *Environ. Int.* **94**, 76–82. <https://doi.org/10.1016/j.envint.2016.05.004>.
- Cienciewicki, J., and Jaspers, I. (2007). Air pollution and respiratory viral infection. *Inhal. Toxicol.* **19**, 1135–1146. <https://doi.org/10.1080/08958370701665434>.
- Wang, Y., Wang, T., Xu, M., et al. (2020). Independent effect of main components in particulate matter on DNA methylation and DNA methyltransferase: a molecular epidemiology study. *Environ. Int.* **134**, 105296. <https://doi.org/10.1016/j.envint.2019.105296>.
- Kan, H.D., Chen, B.H., Fu, C.F., et al. (2005). Relationship between ambient air pollution and daily mortality of SARS in Beijing. *Biomed. Environ. Sci.* **18**, 1–4.
- Pansini, R., and Fornacca, D. (2020). COVID-19 higher mortality in Chinese regions with chronic exposure to lower air quality. *Front. Public Health* **8**, 597753. <https://doi.org/10.3389/fpubh.2020.597753>.
- Wu, X., Nethery, R.C., Sabath, B.M., et al. (2020). Exposure to air pollution and COVID-19 mortality in the United States: a nationwide cross-sectional study. *medRxiv*. . 2020. 2004.2005.20054502. <https://doi.org/10.1101/2020.04.05.20054502>.
- Yao, Y., Pan, J., Liu, Z., et al. (2020). No Association of COVID-19 transmission with temperature or UV radiation in Chinese cities. *Eur. Respir. J.* 2000517. <https://doi.org/10.1183/13993003.00517-2020>.
- Notari, A. (2021). Temperature dependence of COVID-19 transmission. *Sci. Total Environ.* **763**, 144390. <https://doi.org/10.1016/j.scitotenv.2020.144390>.
- Ma, Y., Zhao, Y., Liu, J., et al. (2020). Effects of temperature variation and humidity on the death of COVID-19 in Wuhan, China. *Sci. Total Environ.* **724**, 138226. <https://doi.org/10.1016/j.scitotenv.2020.138226>.
- Chen, R., Yin, P., Wang, L., et al. (2018). Association between ambient temperature and mortality risk and burden: time series study in 272 main Chinese cities. *BMJ* **363**, k4306. <https://doi.org/10.1136/bmj.k4306>.
- Martens, D.S., Plusquin, M., Cox, B., et al. (2019). Early biological aging and fetal exposure to high and low ambient temperature: a birth cohort study. *Environ. Health Perspect.* **127**, 117001. <https://doi.org/10.1289/EHP5153>.
- Lam, H.C.Y., Chan, J.C.N., Luk, A.O.Y., et al. (2018). Short-term association between ambient temperature and acute myocardial infarction hospitalizations for diabetes mellitus patients: a time series study. *PLoS Med.* **15**, e1002612. <https://doi.org/10.1371/journal.pmed.1002612>.
- Chen, S., Zhang, Z., Yang, J., et al. (2020). Fangcang shelter hospitals: a novel concept for responding to public health emergencies. *Lancet* **395**, 1305–1314. [https://doi.org/10.1016/S0140-6736\(20\)30744-3](https://doi.org/10.1016/S0140-6736(20)30744-3).

32. Chen, S.Y., Chan, C.C., and Su, T.C. (2017). Particulate and gaseous pollutants on inflammation, thrombosis, and autonomic imbalance in subjects at risk for cardiovascular disease. *Environ. Pollut.* **223**, 403–408. <https://doi.org/10.1016/j.envpol.2017.01.037>.
33. Hassanvand, M.S., Naddafi, K., Kashani, H., et al. (2017). Short-term effects of particle size fractions on circulating biomarkers of inflammation in a panel of elderly subjects and healthy young adults. *Environ. Pollut.* **223**, 695–704. <https://doi.org/10.1016/j.envpol.2017.02.005>.
34. Hajat, A., Allison, M., Diez-Roux, A.V., et al. (2015). Long-term exposure to air pollution and markers of inflammation, coagulation, and endothelial activation: a repeat-measures analysis in the Multi-Ethnic Study of Atherosclerosis (MESA). *Epidemiology* **26**, 310–320. <https://doi.org/10.1097/EDE.0000000000000267>.
35. Li, D., Chen, Y., Liu, H., et al. (2020). Immune dysfunction leads to mortality and organ injury in patients with COVID-19 in China: insights from ERS-COVID-19 study. *Signal. Transduct. Target Ther.* **5**, 62. <https://doi.org/10.1038/s41392-020-0163-5>.
36. Adar, S.D., Filigrana, P.A., Clements, N., et al. (2014). Ambient coarse particulate matter and human health: a systematic review and meta-analysis. *Curr. Environ. Health Rep.* **1**, 258–274. <https://doi.org/10.1007/s40572-014-0022-z>.
37. Cai, J., Meng, X., Wang, C., et al. (2016). The cold effects on circulatory inflammation, thrombosis and vasoconstriction in type 2 diabetic patients. *Sci. Total Environ.* **568**, 271–277. <https://doi.org/10.1016/j.scitotenv.2016.06.030>.
38. Zhou, F., Yu, T., Du, R., et al. (2020). Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* **395**, 1054–1062. [https://doi.org/10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3).
39. Immovilli, P., Morelli, N., Antonucci, E., et al. (2020). COVID-19 mortality and ICU admission: the Italian experience. *Crit. Care* **24**, 228. <https://doi.org/10.1186/s13054-020-02957-9>.
40. Wang, J., Saguner, A.M., An, J., et al. (2020). Dysfunctional coagulation in COVID-19: from cell to bedside. *Adv. Ther.* **37**, 3033–3039. <https://doi.org/10.1007/s12325-020-01399-7>.
41. Mehta, P., McAuley, D.F., Brown, M., et al. (2020). COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* **395**, 1033–1034. [https://doi.org/10.1016/S0140-6736\(20\)30628-0](https://doi.org/10.1016/S0140-6736(20)30628-0).
42. Schwartz, J., Sarnat, J.A., Coull, B.A., et al. (2007). Effects of exposure measurement error on particle matter epidemiology: a simulation using data from a panel study in Baltimore, MD. *J. Expo. Sci. Environ. Epidemiol.* **17** (Suppl 2), S2–S10. <https://doi.org/10.1038/sj.jes.7500619>.
43. VanderWeele, T.J. (2019). Principles of confounder selection. *Eur. J. Epidemiol.* **34**, 211–219. <https://doi.org/10.1007/s10654-019-00494-6>.
44. Zhao, B.T., Su, Y.X., He, S.S., et al. (2016). Evolution and comparative assessment of ambient air quality standards in China. *J. Integr. Environ. Sci.* **13**, 85–102. <https://doi.org/10.1080/1943815x.2016.1150301>.
45. Qi, J., Ruan, Z., Qian, Z.M., et al. (2020). Potential gains in life expectancy by attaining daily ambient fine particulate matter pollution standards in mainland China: a modeling study based on nationwide data. *PLoS Med.* **17**, e1003027. <https://doi.org/10.1371/journal.pmed.1003027>.
46. Liang, Z., Yang, Y., Qian, Z., et al. (2019). Ambient PM2.5 and birth outcomes: estimating the association and attributable risk using a birth cohort study in nine Chinese cities. *Environ. Int.* **126**, 329–335. <https://doi.org/10.1016/j.envint.2019.02.017>.
47. Altman, D.G., and Bland, J.M. (2003). Statistics Notes - interaction revisited: the difference between two estimates. *BMJ* **326**, 219. <https://doi.org/10.1136/bmj.326.7382.219>.
48. Curtin, F., and Schulz, P. (1998). Multiple correlations and Bonferroni's correction. *Biol. Psychiatry* **44**, 775–777. [https://doi.org/10.1016/s0006-3223\(98\)00043-2](https://doi.org/10.1016/s0006-3223(98)00043-2).

## ACKNOWLEDGMENTS

We would like to extend our sincere gratitude and admiration to academician Jianguo Xu at State Key Laboratory of Infectious Disease Prevention and Control, National Institute for Communicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, for his guidance, support, numerous helpful suggestions and continuous encouragement on this study. We also thank all the staff for their comments on a draft version of the manuscript. This study was supported by the National Natural Science Foundation of China (82041021 and 42041001), the Bill & Melinda Gates Foundation (INV-006371), and the General Program of the State Key Laboratory of Infectious Disease Prevention and Control of China (2020SKLID201).

## AUTHOR CONTRIBUTIONS

H.L., and Q.L. had the idea for and designed the study and had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. F.T. and X.L. drafted the paper. F.T., Z.Q., and A.L. contributed to the exposure assessment. F.T., H.L., S.Z., Y.N., and L.Q. did the analysis. X.L., Y.N., L.Q., Q.C., and S.Z. collected the data. All authors critically revised the manuscript for important intellectual content and gave final approval for the version to be published.

## DECLARATION OF INTERESTS

The authors declare they have no competing interests.

## SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.xinn.2021.100139>.

## LEAD CONTACT WEBSITE

The website of the lead contact is at <http://sph.sysu.edu.cn/teacher/361>.



**The Innovation, Volume 2**

**Supplemental Information**

**Ambient air pollution and low temperature**

**associated with case fatality of COVID-19:**

**A nationwide retrospective cohort study in China**

**Fei Tian, Xiaobo Liu, Qingchen Chao, Zhengmin (Min) Qian, Siqi Zhang, Li Qi, Yanlin Niu, Lauren D. Arnold, Shiyu Zhang, Huan Li, Hualiang Lin, and Qiyong Liu**

## Supplementary Materials

### Contents

Table S1. The univariate analysis of risk factors for COVID-19 fatality in China.

Table S2. Characteristics of study participants by survival status in China using resampling approach.

Table S3. Spearman's correlation coefficients among meteorological variables and air pollutants during the different exposure windows.

Table S4. Sensitivity analyses for the hazard ratios of COVID-19 fatality associated with each 10  $\mu\text{g}/\text{m}^3$  increase in  $\text{PM}_{2.5}$ ,  $\text{PM}_{10}$ ,  $\text{SO}_2$ ,  $\text{NO}_2$ , and  $\text{O}_3$  in different model parameters.

Table S5. Hazard ratios of COVID-19 fatality mortality associated with 10  $\mu\text{g}/\text{m}^3$  increase in  $\text{PM}_{2.5}$ ,  $\text{PM}_{10}$ ,  $\text{NO}_2$ , and  $\text{O}_3$  in the two-pollutant models.

Table S6. Hazard ratios and 95% CI of case fatality of COVID-19 associated with environmental factors of overall exposure windows in China.

Figure S1. The distribution of the COVID-19 case fatality rate in China.

Figure S2. The distribution of the average temperature during the COVID-19 epidemic in China.

Figure S3. The distribution of the average concentration of  $\text{PM}_{2.5}$  during the COVID-19 epidemic in China.

Figure S4. The distribution of the average concentration of  $\text{PM}_{10}$  during the COVID-19 epidemic in China.

Figure S5. The distribution of the average concentration of  $\text{NO}_2$  during the COVID-19 epidemic in China.

Figure S6. The distribution of the average concentration of  $\text{SO}_2$  during the COVID-19 epidemic in

China.

Figure S7. The distribution of the average concentration of O<sup>3</sup> during the COVID-19 epidemic in

China.

Figure S8. Hazard ratios (HRs) and 95% CIs of COVID-19 case-fatality associated with each 10 µg/m<sup>3</sup> increase in air pollutant concentrations at different lags in China.

Effect estimation using the date of symptom onset (A) and the date of diagnosis (B) are shown.

Figure S9. Hazard ratios (HRs) and 95% CIs of case-fatality of COVID-19 associated with each 1 °C decrease in temperature at different lag days in China. Effect estimations using the date of onset and diagnosis are shown.

Text S1. Regression equations for Cox regression models for ambient air pollution.

Text S2. Regression equations for Cox regression models for ambient temperature.

Table S1. The univariate analysis of risk factors for COVID-19 fatality in China.

Variables	Univariate Analysis						
	Coefficient	SE	HR (95% CI)	Likelihood-ratio	Wald	Score (log rank)	P Value *
<b>Age group</b>				25624	21357	31939	
< 25 years			1.00 (Ref)				
25-65 years	2.57	0.17	1.87 (1.26, 2.76)				<0.01
≥ 65 years	4.64	0.17	5.64 (3.88, 8.23)				<0.01
<b>Sex</b>				1853	1780	1829	
Male			1.00 (Ref)				
Female	-0.57	0.01	0.56 (0.54, 0.58)				<0.01
<b>Occupation</b>				8072	5111	6507	
Medical-related			1.00 (Ref)				
Service-related	0.09	0.14	1.10 (0.83, 1.44)				0.51
Office worker	0.59	0.09	1.82 (1.52, 2.18)				<0.01
Home worker	2.30	0.09	10.04 (8.42, 11.96)				<0.01
Others	1.75	0.09	2.91 (1.59, 5.35)				<0.01
<b>Residence</b>				627.9	649.5	655.9	
Local			1.00 (Ref)				
Migrant	0.34	0.01	1.41 (1.37, 1.45)				<0.01
<b>Severity</b>				32715	39291	72970	
Mild/asymptomatic			1.00 (Ref)				
Moderate	-0.30	0.02	0.73 (0.71, 0.77)				0.89
Severe	1.44	0.01	4.24 (4.09, 4.39)				< 0.01

Critical	3.02	0.02	20.47 (19.74, 21.23)				< 0.01
<b>Hospital transfer</b>				15.62	15.48	15.48	
No			1.00 (Ref)				
Yes	-0.06	0.01	0.94 (0.91, 0.97)				< 0.01
<b>Location</b>				6333	5126	5756	
Non-Wuhan			1.00 (Ref)				
Wuhan	1.19	0.02	3.27 (3.17, 3.38)				< 0.01
<b>Province</b>				6119	2941	4225	
Inside Hubei			1.00 (Ref)				
Outside Hubei	-2.12	0.04	0.12 (0.11, 0.13)				< 0.01
<b>Lockdown</b> †				1270	1396	1432	
Before			1.00 (Ref)				
After	-0.55	0.01	0.57 (0.56, 0.59)				< 0.01
<b>GDP Per Capita</b>	-0.06	0.005	0.94 (0.93, 0.95)	115	109.5	109	< 0.01
<b>Hospital Beds per 1000 Persons</b>	1.13	0.03	3.11 (2.95, 3.28)	2691	1729	1852	< 0.01

Note: HR, hazard ratio; GDP, gross domestic product.

\* *P* values were calculated by  $\chi^2$  test or Fisher's exact test, as appropriate.

† Lockdown: On 23 January 2020, the central government of China imposed a lockdown in Wuhan and other cities in Hubei.

Table S2. Characteristics of study participants by survival status in China using resampling approach.

Variable	Deceased (n=3,934)	Alive (n=3,934) §		P-value †
		Resampling value	95% CI	
<b>Environmental factors</b>				
<b>Exposure window I *</b>				
Temperature (°C)	5.90 ± 2.24	6.27 ± 0.06	6.17, 6.37	<0.001
Relative humidity (%)	80.13 ± 5.75	78.09 ± 0.13	77.87, 78.31	<0.001
PM <sub>2.5</sub> (µg/m <sup>3</sup> )	53.71 ± 15.52	51.03 ± 0.03	50.50, 51.57	<0.001
PM <sub>10</sub> (µg/m <sup>3</sup> )	62.02 ± 16.93	59.76 ± 0.04	59.18, 60.42	<0.001
O <sub>3</sub> (µg/m <sup>3</sup> )	54.31 ± 9.03	56.02 ± 0.01	55.77, 56.27	<0.001
SO <sub>2</sub> (µg/m <sup>3</sup> )	7.33 ± 2.11	7.68 ± 0.01	7.58, 7.77	<0.001
NO <sub>2</sub> (µg/m <sup>3</sup> )	22.88 ± 6.61	19.81 ± 0.01	19.63, 19.99	<0.001
<b>Exposure window II *</b>				
Temperature (°C)	7.41 ± 2.76	6.27 ± 0.06	6.18, 6.36	<0.001
Relative humidity (%)	79.66 ± 6.53	78.20 ± 0.13	78.00, 78.41	<0.001
PM <sub>2.5</sub> (µg/m <sup>3</sup> )	45.11 ± 17.32	51.11 ± 0.03	50.58, 51.69	<0.001
PM <sub>10</sub> (µg/m <sup>3</sup> )	49.85 ± 18.66	59.93 ± 0.04	59.21, 60.46	<0.001
O <sub>3</sub> (µg/m <sup>3</sup> )	54.73 ± 9.01	55.97 ± 0.02	55.70, 56.22	<0.001
SO <sub>2</sub> (µg/m <sup>3</sup> )	7.41 ± 2.10	7.65 ± 0.01	7.57, 7.74	<0.001
NO <sub>2</sub> (µg/m <sup>3</sup> )	19.54 ± 6.17	19.91 ± 0.01	19.74, 20.08	<0.001
<b>Age group (n, %)</b>				<0.001
< 25 years	6 (0.15)	203 (5.16)	179, 226	
25-64 years	1146 (29.13)	2905 (73.84)	2860, 2949	
≥ 65 years	2782 (70.72)	826 (21.00)	785, 867	
<b>Sex (n, %)</b>				<0.001
Male	2515 (63.93)	1983 (50.41)	1932, 2033	
Female	1419 (36.07)	1951 (49.59)	1899, 2000	
<b>Occupation (n, %)</b>				<0.001
Medical-related	21 (0.53)	152 (3.86)	132, 171	
Service-related	15 (0.38)	99 (2.52)	83, 115	
Office worker	226 (5.74)	895 (22.75)	849, 937	
Homemaker	3226 (82.00)	2238 (56.89)	2189, 2290	
Others	446 (11.34)	550 (13.98)	515, 588	

<b>Residence (n, %)</b>			<0.001
Permanent	2476 (62.94)	2785 (70.79)	2738, 2833
Temporarily	1458 (37.06)	1149 (29.21)	1101, 1196
<b>Severity (n, %)</b>			<0.001
Mild	806 (20.49)	1627 (41.36)	1576, 1678
Moderate	597 (15.18)	1635 (41.56)	1585, 1683
Severe	1285 (32.66)	585 (14.87)	548, 619
Critical	1246 (31.67)	87 (2.21)	72, 103
<b>Hospital transfer</b>			0.092
No	2927 (74.40)	2886 (73.36)	2839, 2928
Yes	1007 (25.60)	1048 (26.64)	1006, 1095

Note: PM<sub>2.5</sub>, particulate matter with an aerodynamic diameter  $\leq 2.5 \mu\text{m}$ ; PM<sub>10</sub>, particulate matter with an aerodynamic diameter  $\leq 10 \mu\text{m}$ ; SO<sub>2</sub>, sulfur dioxide; NO<sub>2</sub>, nitrogen dioxide; O<sub>3</sub>, ozone.

\* Exposure window I represents the mean exposure value from the date of symptom onset to the date of diagnosis; Exposure window II represents the mean exposure value from the date of diagnosis to the date of death or the end of the study.

† *P*-value were calculated by  $\chi^2$  test or Fisher's exact test, as appropriate.

§ The results of the alive were based on the 1000 times random resampling.

Table S3. Spearman's correlation coefficients among meteorological variables and air pollutants during the different exposure windows. \*

Variables	Temperature	Relative humidity	PM <sub>2.5</sub>	PM <sub>10</sub>	O <sub>3</sub>	NO <sub>2</sub>	SO <sub>2</sub>
<b>Exposure I †</b>							
Temperature	1.00						
Relative humidity	-0.14	1.00					
PM <sub>2.5</sub>	0.36	-0.16	1.00				
PM <sub>10</sub>	0.35	-0.17	0.96	1.00			
NO <sub>2</sub>	0.19	0.02	0.45	0.43	1.00		
SO <sub>2</sub>	0.43	-0.39	0.34	0.38	0.31	1.00	
O <sub>3</sub>	0.01	-0.59	0.23	0.22	-0.21	0.18	1.00
<b>Exposure II †</b>							
Temperature	1.00						
Relative humidity	-0.34	1.00					
PM <sub>2.5</sub>	0.45	-0.30	1.00				
PM <sub>10</sub>	0.29	-0.58	0.76	1.00			
NO <sub>2</sub>	-0.12	-0.13	0.15	0.23	1.00		
SO <sub>2</sub>	0.16	-0.25	0.28	0.37	0.31	1.00	
O <sub>3</sub>	-0.06	-0.28	0.01	0.14	-0.16	0.08	1.00

Note: PM<sub>2.5</sub>, particulate matter with an aerodynamic diameter  $\leq 2.5$   $\mu\text{m}$ ; PM<sub>10</sub>, particulate matter with an aerodynamic diameter  $\leq 10$   $\mu\text{m}$ ; SO<sub>2</sub>, sulfur dioxide; NO<sub>2</sub>, nitrogen dioxide; O<sub>3</sub>, ozone.

\* All pairwise correlation coefficients were statistically significant ( $P < 0.05$ ).

† Exposure window I represents the mean exposure value from the date of symptom onset to the date of diagnosis; Exposure window II represents the mean exposure value from the date of diagnosis to the date of death or the end of the study.



Table S4. Sensitivity analyses for the hazard ratios of COVID-19 fatality associated with each 10 µg/m<sup>3</sup> increase in PM<sub>2.5</sub>, PM<sub>10</sub>, SO<sub>2</sub>, NO<sub>2</sub>, and O<sub>3</sub> in different model parameters.

<b>Models</b> †	<b>PM<sub>2.5</sub></b> *	<b>PM<sub>10</sub></b> *	<b>NO<sub>2</sub></b> *	<b>SO<sub>2</sub></b> *	<b>O<sub>3</sub></b> *
df=2 for temperature	1.11 (1.09, 1.13)	1.10 (1.08, 1.12)	1.25 (1.16, 1.34)	1.10 (0.95, 1.27)	1.08 (1.02, 1.14)
df=4 for temperature	1.10 (1.07, 1.13)	1.10 (1.07, 1.12)	1.27 (1.19, 1.35)	1.10 (0.95, 1.26)	1.09 (1.03, 1.15)
df=5 for relative humidity	1.11 (1.08, 1.13)	1.10 (1.08, 1.12)	1.27 (1.18, 1.35)	1.11 (0.95, 1.27)	1.09 (1.03, 1.14)
df=2 for relative humidity	1.10 (1.08, 1.13)	1.10 (1.08, 1.13)	1.26 (1.19, 1.36)	1.10 (0.95, 1.27)	1.12 (1.03, 1.20)
df=4 for relative humidity	1.11 (1.08, 1.14)	1.10 (1.08, 1.12)	1.27 (1.19, 1.35)	1.09 (0.95, 1.25)	1.09 (1.03, 1.14)
df=5 for relative humidity	1.09 (1.07, 1.12)	1.10 (1.08, 1.13)	1.26 (1.18, 1.36)	1.10 (0.95, 1.27)	1.09 (1.03, 1.14)
df=6 for relative humidity	1.11 (1.08, 1.14)	1.10 (1.08, 1.12)	1.27 (1.19, 1.35)	1.10 (0.96, 1.27)	1.10 (1.05, 1.15)

\*Effect estimated using Window I, and exposure window I represents the mean exposure value from the date of symptom onset to the date of diagnosis.

† Multivariate model, adjusted for age, sex, occupation, residence, severity of the illness, location, and transfer history, temporal trend, lockdown, city-level GDP, hospital beds per 1000 persons, temperature (only for the pollutants) and relative humidity.

Table S5. Hazard ratios of COVID-19 fatality mortality associated with 10  $\mu\text{g}/\text{m}^3$  increase in  $\text{PM}_{2.5}$ ,

$\text{PM}_{10}$ ,  $\text{NO}_2$ , and  $\text{O}_3$  in the two-pollutant models.

<b>Pollutants</b>	<b>Models</b>	<b>Hazard Ratios</b>	<b><i>P</i>-value<sup>†</sup></b>
<b><math>\text{PM}_{2.5}</math></b>	Single pollutant model	1.11 (1.09, 1.13)	
	<b>Two- pollutant model</b>		
	Adjust for $\text{O}_3$	1.11 (1.09, 1.14)	0.99
	Adjust for $\text{NO}_2$	1.10 (1.07, 1.13)	0.58
	Adjust for $\text{SO}_2$	1.13 (1.11, 1.16)	0.22
<b><math>\text{PM}_{10}</math></b>	Single pollutant model	1.10 (1.08, 1.13)	
	<b>Two- pollutant model</b>		
	Adjust for $\text{O}_3$	1.10 (1.08, 1.13)	0.99
	Adjust for $\text{NO}_2$	1.09 (1.07, 1.12)	0.58
	Adjust for $\text{SO}_2$	1.13 (1.10, 1.15)	0.09
<b><math>\text{NO}_2</math></b>	Single pollutant model	1.27 (1.19, 1.35)	
	<b>Two- pollutant model</b>		
	Adjust for $\text{PM}_{2.5}$	1.08 (0.99, 1.17)	<0.01
	Adjust for $\text{PM}_{10}$	1.06 (0.98, 1.16)	<0.01
	Adjust for $\text{O}_3$	1.28 (1.19, 1.36)	0.86
	Adjust for $\text{SO}_2$	1.32 (1.23, 1.43)	0.44
<b><math>\text{O}_3</math></b>	Single pollutant model	1.09 (1.03, 1.14)	
	<b>Two- pollutant model</b>		
	Adjust for $\text{PM}_{2.5}$	1.01 (0.95, 1.06)	0.04
	Adjust for $\text{PM}_{10}$	1.01 (0.96, 1.06)	0.04
	Adjust for $\text{NO}_2$	1.09 (1.04, 1.15)	0.99
	Adjust for $\text{SO}_2$	1.09 (1.04, 1.14)	0.99

\*Effect estimated using Window I.

<sup>†</sup> Estimated using likelihood ratio test by comparing the single-pollutant model and each nested two-pollutant model.

Table S6. Hazard ratios and 95% CI of case fatality of COVID-19 associated with environmental factors of overall exposure windows in China \*.

Variables	Hazard Ratios (95% CI)	
	Crude §	Adjusted ‡
Temperature	1.30 (1.28, 1.32)	2.34 (2.30, 2.39)
PM <sub>2.5</sub>	1.22 (1.14, 1.30)	1.31 (1.26, 1.37)
PM <sub>10</sub>	1.20 (1.18, 1.22)	1.29 (1.15, 1.44)
NO <sub>2</sub>	1.32 (1.13, 1.51)	1.52 (1.23, 1.80)
SO <sub>2</sub>	0.75 (0.52, 0.99)	1.06 (0.83, 1.29)
O <sub>3</sub>	0.93 (0.73, 1.11)	1.32 (1.14, 1.51)

Note: PM<sub>2.5</sub>, particulate matter with an aerodynamic diameter  $\leq 2.5 \mu\text{m}$ ; PM<sub>10</sub>, particulate matter with an aerodynamic diameter  $\leq 10 \mu\text{m}$ ; SO<sub>2</sub>, sulfur dioxide; NO<sub>2</sub>, nitrogen dioxide; O<sub>3</sub>, ozone.

\* Exposure window represents the exposure window of whole infection course (from the symptom onset to death/recovery or end of the study)

§ Crude model, with no adjustment.

‡ Multivariate model, adjusted for age, sex, occupation, residence, severity of the illness, location, and transfer history, temporal trend, lockdown, city-level GDP, hospital beds per 1000 persons, temperature (only for the pollutants) and relative humidity.




Figure S1. The distribution of fatality rates across the 321 cities in China.




Figure S2. The distribution of the average temperature during the COVID-19 epidemic in China.




Figure S3. The distribution of the average concentration of PM<sub>2.5</sub> during the COVID-19 epidemic in China.




Figure S4. The distribution of the average concentration of PM<sub>10</sub> during the COVID-19 epidemic in China.




Figure S5. The distribution of the average concentration of NO<sub>2</sub> during the COVID-19 epidemic in China.






Figure S6. The distribution of the average concentration of SO<sub>2</sub> during the COVID-19 epidemic in China.




Figure S7. The distribution of the average concentration of O<sub>3</sub> during the COVID-19 epidemic in China.




Figure S8. Hazard ratios (HRs) and 95% CIs of COVID-19 case-fatality associated with each 10 µg/m<sup>3</sup> increase in air pollutant concentrations at different lags in China.

Effect estimation using the date of symptom onset (A) and the date of diagnosis (B) are shown.




Figure S9. Hazard ratios (HRs) and 95% CIs of case-fatality of COVID-19 associated with each 1 °C decrease in temperature at different lag days in China. Effect estimations using the date of onset and diagnosis are shown.

Text S1. Regression equations for Cox regression model III for ambient air pollution.

$$h(t) = h_0 * \exp(\beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_4 X_4 + \beta_5 X_5 + \beta_6 X_6 + \beta_7 X_7 + \beta_8 X_8 + \beta_9 X_9 + \beta_{10} X_{10} \\ + \beta_{11} X_{11} + \beta_{12} X_{12} + \beta_{13} X_{13} + \beta_{14} X_{14})$$

where:

$t$  represents the survival time;

$h(t)$  is the hazard function determined by a set of variates ( $X_1, X_2, \dots, X_{14}$ );

The term  $h_0$  is called the baseline hazard. It corresponds to the value of the hazard if all the  $X_i$  are equal to zero. The 't' in  $h(t)$  reminds us that the hazard may vary over time.

The coefficients ( $\beta_1, \beta_2, \dots, \beta_{14}$ ) = effects parameter estimates;

$X_1$  = gaseous pollutants (PM<sub>2.5</sub>, PM<sub>10</sub>, NO<sub>2</sub>, SO<sub>2</sub>, and O<sub>3</sub>);

$X_2$  = sex (female or male);

$X_3$  = age group (<25 years, 25-65 years, and ≥65 years);

$X_4$  = residence (local or migrant);

$X_5$  = occupation (medical-related, service-related, office worker, home worker, and others);

$X_6$  = severity of the illness (mild/asymptomatic, moderate, severe, critical);

$X_7$  = location (Wuhan or Non-Wuhan);

$X_8$  = transfer to a better hospital or not (yes or no);

$X_9$  = time trend;

$X_{10}$  = temperature;

$X_{11}$  = relative humidity;

$X_{12}$  = lockdown (before or after);

$X_{13}$  = city-level GDP;

$X_{14}$  = hospital beds per 1000 persons.

Text S2. Regression equations for Cox regression model III for ambient temperature.

$$h(t) = h_0 * \exp(\beta_1 X_t + \beta_2 X_2 + \beta_3 X_3 + \beta_4 X_4 + \beta_5 X_5 + \beta_6 X_6 + \beta_7 X_7 + \beta_8 X_8 + \beta_9 X_9 + \beta_{10} X_{10} + \beta_{11} X_{11} + \beta_{12} X_{12} + \beta_{13} X_{13})$$

where:

$t$  represents the survival time;

$h(t)$  is the hazard function determined by a set of variates ( $X_t, X_2, \dots, X_{13}$ );

The term  $h_0$  is called the baseline hazard. It corresponds to the value of the hazard if all the  $X_i$  are equal to zero. The 't' in  $h(t)$  reminds us that the hazard may vary over time.

The coefficients ( $\beta_1, \beta_2, \dots, \beta_{13}$ ) = effects parameter estimates;

$X_t$  = temperature;

$X_2$  = sex (female or male);

$X_3$  = age group (<25 years, 25-65 years, and  $\geq 65$  years);

$X_4$  = residence (local or migrant);

$X_5$  = occupation (medical-related, service-related, office worker, home worker, and others);

$X_6$  = severity of the illness (mild/asymptomatic, moderate, severe, critical);

$X_7$  = location (Wuhan or Non-Wuhan);

$X_8$  = transfer to a better hospital or not (yes or no);

$X_9$  = time trend;

$X_{10}$  = relative humidity;

$X_{11}$  = lockdown (before or after);

$X_{12}$  = city-level GDP;

$X_{13}$  = hospital beds per 1000 persons.