

**Supplemental Table 2: Baseline characteristics of young CRVO patients who had hypercoagulable testing or progressed to being diagnosed with a hypercoagulable state at 1 year**

	Hypercoagulable testing				Hypercoagulable State at 1 year			
	No (N=731)	Yes (N=450)	Total (N=1181)	p value	No (N=1045)	Yes (N=136)	Total (N=1181)	p value
<b>Gender (F)</b>	307 (42.0%)	204 (45.3%)	511 (43.3%)	0.261	449 (43.0%)	62 (45.6%)	511 (43.3%)	0.562
<b>Age (Mean (SD))</b>	40.7 (8.590)	39.1 (8.707)	40.1 (8.662)	0.004	40.5 (8.395)	37.2 (10.071)	40.1 (8.662)	<0.001
<b>Smoking Hx</b>	110 (15.0%)	46 (10.2%)	156 (13.2%)	0.017	143 (13.7%)	13 (9.6%)	156 (13.2%)	0.181
<b>Hypertension Hx</b>	298 (40.8%)	141 (31.3%)	439 (37.2%)	0.001	395 (37.8%)	44 (32.4%)	439 (37.2%)	0.216
<b>Diabetes Mellitus Hx</b>	156 (21.3%)	60 (13.3%)	216 (18.3%)	<0.001	197 (18.9%)	19 (14.0%)	216 (18.3%)	0.166
<b>Hyperlipidemia Hx</b>	294 (40.2%)	168 (37.3%)	462 (39.1%)	0.324	412 (39.4%)	50 (36.8%)	462 (39.1%)	0.550
<b>Hypercoagulable state at 1 year</b>	43 (5.9%)	93 (20.7%)	136 (11.5%)	<0.001	--	--	--	