Supplementary Online Content

Ferm ML, DeSalvo DJ, Prichett LM, Sickler JK, Wolf RM, Channa R. Clinical and demographic factors associated with diabetic retinopathy among young patients with diabetes. *JAMA Netw Open.* 2021;4(9):e2126126. doi:10.1001/jamanetworkopen.2021.26126

eTable. Multivariable Regression Analysis of Patients With Diabetic Retinopathy as Outcome Excluding Insurance

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Multivariable Regression Analysis of Patients With Diabetic Retinopathy as Outcome Excluding Insurance

T1D (n=1207)	Odds Ratio (95% Confidence Interval)	Standard Error	Z	P> z
Duration of Diabetes	1.172 (1.093-1.256)	0.041	4.480	<0.001
	1.172 (1.093-1.230)	0.041	4.460	<0.001
<u>Race (categorical)</u>				
White (NH)	Reference			
Black/African				
American (NH)	1.843 (0.867-3.918)	0.709	1.590	0.11
Hispanic/Other	0.635 (0.266-1.517)	0.282	-1.020	0.31
Mean HbA1c	1.168 (0.996-1.369)	0.095	1.910	0.06
Pump (y)	0.440 (0.204-0.948)	0.172	-2.100	0.04

	Odds Ratio (95%	Standard		
T2D (n=376)	Confidence Interval)	Error	Z	P> z
Duration of Diabetes	1.259 (1.076-1.474)	0.101	2.870	0.004
Race (categorical)				
White (NH)	Reference			
Black/African				
American (NH)	0.927 (0.274-3.138)	0.577	-0.120	0.90
Hispanic/Other	1.000	(omitted)		
Mean HbA1c	1.035 (0.837-1.280)	0.112	0.320	0.75

T1D = type 1 diabetes, T2D = type 2 diabetes, DR = diabetic retinopathy, NH = non-Hispanic, HbA1c = hemoglobin A1c