

Appendix S1. The pediRES-Q Collaborative Investigators: Kamal Abulebda, Riley Hospital for Children, Indianapolis, IN, US; Diane Atkins, University of Iowa Stead Family Children's Hospital, Iowa City, IA, US; Shilpa Balikai, University of Iowa Stead Family Children's Hospital, Iowa City, IA, US; Marc Berg, Lucile Packard Children's Hospital, Palo Alto, CA, US; Robert Berg, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Utpal Bhalala, Children's Hospital of San Antonio, San Antonio, TX, US; Matthew S. Braga, Dartmouth-Hitchcock Medical Center, Lebanon, NH, US; Corinne Buysse, Erasmus MC-Sophia Children's Hospital, Rotterdam, NL; Corrado Cecchetti, Ospedale Pediatrico Bambino Gesù, Rome, IT; Adam Chenq, Alberta Children's Hospital, Calgary, AB, CA; Andrea Christoff, Children's Hospital at Westmead, Sydney, NSW; Kelly Corbett, Dartmouth-Hitchcock Medical Center, Lebanon, NH, US; Allan DeCaen, Stollery Children's Hospital, Edmonton, AB, CA; Gabry deJong, Erasmus MC–Sophia Children's Hospital, Rotterdam, NL; Jimena del Castillo, Hospital Maternoinfantil Gregorio Marañón, Madrid, ES; Maya Dewan, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, US; Aaron Donoghue, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Jordan Duval-Arnould, Johns Hopkins University School of Medicine, Baltimore, MD, US; Ivie Esangbedo, UT Southwestern Dallas Children's Medical Center, Dallas, TX, US; Michael Flaherty, Massachusetts General Hospital, Boston, MA, US; Stuart Friess, St. Louis Children's Hospital, St. Louis, MO, US; Sandeep Gangadharan, Mount Sinai Kravis Children's Hospital, New York, NY, US; Orsola Gawronski, Ospedale Pediatrico Bambino Gesù, Rome, IT; Jonathan Gilleland, Alberta Children's Hospital, Calgary, AB, CA; Heather Griffis, Healthcare Analytics Unit, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Richard Hanna, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Helen Harvey, Rady Children's Hospital, San Diego, CA, US; Ilana Harwayne-Gidansky, Stony Brook Hospital, Stony Brook, NY, US; Sarah Haskell, University of Iowa Stead Family Children's Hospital, Iowa City, IA, US; Jennifer Hayes, Children's Hospital of Orange County, Los Angeles, CA, US; Kiran Heber, Children's Healthcare of Atlanta, Atlanta, GA, US; Betsy Hunt, Johns Hopkins University School of Medicine, Baltimore, MD, US; Takanari Ikeyama, Aichi Children's Health and Medical Center, Obu, Aichi, JP; Priti Jani, The University of Chicago Medicine Comer Children's Hospital, Chicago, IL, US; Kaitlin Jones, Children's Healthcare of Atlanta, Atlanta, GA, US; Monica Kleinman, Boston Children's Hospital, Boston, MA, US; Lynda Knight, Lucile Packard Children's Hospital Stanford, Palo Alto, CA, US; Hiroshi Kurosawa, Hyogo Prefectural Kobe Children's Hospital, Kobe, Hyogo, JP; Javier Lasa, Texas Children's Hospital, Houston, TX, US; Kasper Glerup Lauridsen, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Aarhus University Hospital, Aarhus, DK; Tara Lemoine, Valley Children's Hospital, Madera, CA, US; Tensing Maa, Nationwide Children's Hospital, Columbus, OH, US; Dori-Ann Martin, Alberta Children's Hospital, Calgary, AB, CA; Elizabeth Masse, Seattle Children's Hospital, Seattle, WA, US; Luz Marina Mejia, Instituto de Ortopedia Infantil Roosevelt, Bogota, CO; Michael Meyer, Children's Hospital of Wisconsin, Milwaukee, WI, US; Yee Hui Mok, KK Women's & Children's Hospital, SG; Ryan Morgan, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Vinay Nadkarni, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Sholeen Nett, Dartmouth-Hitchcock Medical Center, Lebanon, NH, US; Dana Niles, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Michelle Olson, Children's Hospital of Richmond, Richmond, VA, US; Tara Petersen, Children's Hospital of Wisconsin, Milwaukee, WI, US; Gene Ong, KK Women's & Children's Hospital, SG; Prakad Rajapreyar, Children's Hospital of Wisconsin, Milwaukee, WI, US; Tia Raymond, Medical City Children's Hospital, Dallas, TX, US; Joan Roberts, Seattle Children's Hospital, Seattle, WA, US; Lindsay Ryerson, Stollery Children's Hospital, Edmonton, AB, CA; Anita Sen, NewYork-Presbyterian Morgan Stanley Children's Hospital, New York, NY, US; Marcy Singleton, Dartmouth-Hitchcock Medical Center, Lebanon, NH, US; Sophie Skellet, Great Ormond Street Hospital, London, UK; **Daniel Stromberg**, Dell Children's Medical Center, Austin, TX, US; **Felice Su**, Lucile Packard Children's Hospital Stanford, Palo Alto, CA, US; Robert Sutton, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Todd Sweberg, Cohen Children's Medical Center, New Hyde Park, NY, US; Oscar Tegg, The Children's Hospital at Westmead, Sydney, NSW; Ken Tegtmeyer, Cincinnati Children's Hospital, Cincinnati, OH, US; Alexis Topjian, The Children's Hospital of Philadelphia, Philadelphia, PA, US; Wendy Van Ittersum, Akron Children's Hospital, Akron, OH, US; Javier Urbano Villaescusa; Hospital Maternoinfantil Gregorio Marañón, Madrid, ES; Ichiro Watanabe, Tokyo Metropolitan Children's Hospital, Fuchū, Tokyo, JP; Denise Welsby, Great Ormond Street Hospital, London, UK; Jesse Wenger, Seattle Children's Hospital, Seattle, WA, US; Heather **Wolfe**, The Children's Hospital of Philadelphia, Philadelphia, PA, US; **Andrea Yeo**, National University Children's Medical Institute, SG; Pricilla Yu, UT Southwestern Dallas Children's Medical Center, Dallas, TX, US.

Figure S1. The distribution of defibrillation current (Ohms) for the first shock ( $shock_1$ ) among the 39 subjects with defibrillation for VF (ventricular fibrillation).

