

1 **Supplemental Digital Content**

2 **Table 1. Subject discontinuations and analysis populations**

All subjects (N=111)	Month 1	Month 3	Month 6
ITT population (N=83)	83	71	66
PP population (N=55)	55	55	55
Subjects discontinued, n (%)	28 (25.2)	40 (36.0)	45 (40.5)
Subjects remaining in the study, n (%)	83 (74.8)	71 (64.0)	66 (59.5)

3 *The ITT analysis set consisted of subjects who received at least one dose of vortioxetine, and had pre-*
 4 *intervention and at least one post-intervention assessment of efficacy.*

5 *The PP analysis set was a subset of the ITT analysis set, and consisted of subjects who completed all*
 6 *scheduled visits without major protocol deviations or violations.*

7 *Percentages are based on the total number of enrolled subjects (N=111).*

8 **Supplemental Digital Content**9 **Table 2. Adverse drug reactions**

Summary of ADRs		N = 111
System Organ Class	n (%) of subjects	No. of events
Preferred Term		
Any ADRs	3 (2.7)	4
Serious ADRs		
Not serious		4
Severity		
Mild		4
Changes to study product due to event		
Dose not changed		4
ADRs (diagnosis)		
Nausea	2 (1.8)	2
Gastritis	1 (0.9)	1
Headache	1 (0.9)	1

10 *ADRs were any AEs that were assessed by the investigators to be possibly, probably or definitely related*

11 *to the study treatment*