

Table A1: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Total SHARE population^a in 2004-05 (wave 1) and 2015 (wave 6)

	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
2004-05	50	4057	31.9	22.8 (22.6-23.0)	71.5 (70.8-72.1)	5.7 (5.5-5.9)	17.8 (17.3-18.4)	3.4 (3.3-3.6)	10.7 (10.2-11.2)
	55	4102	27.5	18.8 (18.6-19.0)	68.3 (67.6-69.1)	5.3 (5.1-5.5)	19.3 (18.7-20.0)	3.4 (3.2-3.5)	12.3 (11.8-12.9)
	60	3716	23.2	15.0 (14.8-15.2)	64.7 (63.9-65.6)	4.8 (4.7-5.0)	20.9 (20.1-21.7)	3.3 (3.2-3.5)	14.4 (13.7-15.1)
	65	3345	19.1	11.4 (11.2-11.6)	59.7 (58.7-60.7)	4.4 (4.3-4.6)	23.2 (22.3-24.2)	3.3 (3.1-3.4)	17.1 (16.2-17.9)
	70	2726	15.3	8.1 (7.9-8.3)	52.8 (51.5-54.1)	4.0 (3.8-4.2)	26.4 (25.2-27.6)	3.2 (3.0-3.4)	20.9 (19.7-22.0)
	75	2000	11.8	5.4 (5.2-5.6)	46.0 (44.3-47.7)	3.4 (3.2-3.6)	28.7 (27.1-30.3)	3.0 (2.8-3.2)	25.4 (23.8-26.9)
	80	1183	8.7	3.2 (3.0-3.4)	37.3 (34.9-39.7)	2.7 (2.5-2.9)	31.0 (28.7-33.4)	2.7 (2.5-3.0)	31.7 (29.3-34.0)
	85+	569	6.2	2.1 (1.8-2.3)	33.4 (29.5-37.3)	2.1 (1.8-2.3)	33.0 (29.1-36.9)	2.1 (1.9-2.3)	33.6 (29.7-37.5)
2015	50	3896	33.3	24.0 (23.8-24.1)	72.1 (71.7-72.6)	5.9 (5.7-6.0)	17.7 (17.3-18.1)	3.4 (3.3-3.5)	10.2 (9.9-10.5)
	55	6090	28.8	20.0 (19.9-20.2)	69.6 (69.2-70.1)	5.4 (5.3-5.5)	18.7 (18.3-19.2)	3.3 (3.2-3.5)	11.6 (11.3-12.0)
	60	7046	24.4	16.2 (16.0-16.3)	66.1 (65.6-66.7)	5.0 (4.9-5.1)	20.5 (20.0-21.0)	3.3 (3.2-3.4)	13.4 (13.0-13.8)
	65	7029	20.3	12.5 (12.4-12.6)	61.5 (60.9-62.2)	4.6 (4.5-4.7)	22.5 (21.9-23.1)	3.2 (3.1-3.4)	15.9 (15.4-16.5)
	70	5585	16.4	9.2 (9.0-9.3)	55.7 (54.9-56.5)	4.1 (4.0-4.2)	24.9 (24.1-25.6)	3.2 (3.1-3.3)	19.4 (18.7-20.1)
	75	4430	12.7	6.2 (6.0-6.3)	48.5 (47.5-49.6)	3.4 (3.3-3.6)	27.1 (26.1-28.1)	3.1 (3.0-3.2)	24.4 (23.4-25.3)
	80	2846	9.4	3.7 (3.6-3.8)	39.5 (38.1-40.9)	2.8 (2.7-2.9)	29.9 (28.6-31.3)	2.9 (2.7-3.0)	30.5 (29.2-31.9)
	85+	1937	6.7	2.2 (2.0-2.3)	32.4 (30.3-34.4)	2.1 (2.0-2.3)	31.7 (29.7-33.8)	2.4 (2.3-2.5)	35.9 (33.8-38.0)

^a Denmark, Sweden, Austria, Germany, France, Switzerland, Belgium, Italy, Spain and Greece.

Table A2: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Northern SHARE population^a in 2004-05 (wave 1) and 2015 (Wave 6)

	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
2004-05	50	775	31.3	25.3 (24.9-25.7)	80.7 (79.5-82.0)	4.2 (3.8-4.5)	13.4 (12.3-14.5)	1.8 (1.6-2.1)	5.8 (5.0-6.7)
	55	870	26.8	21.0 (20.6-21.4)	78.3 (76.8-79.7)	4.0 (3.7-4.4)	15.0 (13.7-16.3)	1.8 (1.5-2.1)	6.7 (5.8-7.7)
	60	750	22.6	17.1 (16.7-17.5)	75.7 (74.0-77.4)	3.7 (3.4-4.1)	16.5 (15.0-18.1)	1.8 (1.5-2.0)	7.8 (6.6-8.9)
	65	609	18.5	13.3 (12.9-13.7)	71.8 (69.7-73.8)	3.5 (3.2-3.9)	18.9 (17.0-20.8)	1.7 (1.5-2.0)	9.3 (7.9-10.8)
	70	494	14.8	9.8 (9.4-10.2)	66.0 (63.3-68.6)	3.3 (3.0-3.7)	22.5 (20.0-24.9)	1.7 (1.4-2.0)	11.6 (9.6-13.5)
	75	411	11.5	6.8 (6.3-7.2)	58.9 (55.4-62.5)	3.0 (2.6-3.4)	26.2 (23.0-29.5)	1.7 (1.4-2.0)	14.8 (12.1-17.5)
	80	221	8.5	4.3 (3.8-4.7)	50.3 (45.2-55.4)	2.6 (2.2-3.0)	30.3 (25.5-35.0)	1.6 (1.3-2.0)	19.4 (15.3-23.5)
	85+	152	6.1	2.7 (2.2-3.2)	43.9 (36.0-51.8)	2.1 (1.7-2.6)	35.3 (27.7-42.9)	1.3 (0.9-1.7)	20.8 (14.4-27.3)
2015	50	677	33.0	27.2 (26.9-27.5)	82.5 (81.6-83.4)	3.8 (3.5-4.1)	11.5 (10.7-12.3)	2.0 (1.8-2.2)	6.0 (5.4-6.6)
	55	976	28.4	22.9 (22.6-23.1)	80.4 (79.4-81.4)	3.6 (3.3-3.8)	12.6 (11.7-13.5)	2.0 (1.8-2.2)	7.0 (6.3-7.7)
	60	1224	24.0	18.7 (18.4-19.0)	77.8 (76.6-78.9)	3.4 (3.2-3.6)	14.1 (13.1-15.1)	1.9 (1.7-2.1)	8.1 (7.3-8.9)
	65	1434	19.9	14.8 (14.5-15.1)	74.5 (73.1-75.8)	3.1 (2.9-3.4)	15.7 (14.5-16.9)	2.0 (1.8-2.2)	9.9 (8.8-10.9)
	70	1148	15.9	11.1 (10.8-11.4)	69.8 (68.1-71.6)	2.9 (2.6-3.1)	18.0 (16.4-19.5)	1.9 (1.7-2.2)	12.2 (10.9-13.5)
	75	809	12.3	7.8 (7.5-8.1)	63.4 (61.1-65.8)	2.6 (2.3-2.8)	21.0 (18.9-23.0)	1.9 (1.7-2.1)	15.6 (13.8-17.4)
	80	500	9.1	5.0 (4.7-5.3)	55.7 (52.4-59.0)	2.2 (1.9-2.4)	23.9 (21.0-26.8)	1.9 (1.6-2.1)	20.4 (17.7-23.2)
	85+	377	6.4	3.0 (2.7-3.3)	46.9 (41.9-52.0)	1.7 (1.4-2.0)	26.7 (22.3-31.2)	1.7 (1.4-2.0)	26.3 (21.9-30.8)

^a Denmark and Sweden.

Table A3: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Central SHARE population^a in 2004-05 (Wave 1) and 2015 (Wave 6)

	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
2004-05	50	2093	31.6	24.2 (23.9-24.4)	76.4 (75.5-77.2)	4.8 (4.5-5.0)	15.0 (14.2-15.8)	2.7 (2.5-2.9)	8.6 (7.9-9.3)
	55	2014	27.3	20.0 (19.8-20.3)	73.5 (72.5-74.5)	4.5 (4.2-4.7)	16.5 (15.6-17.4)	2.7 (2.5-2.9)	10.0 (9.2-10.8)
	60	1818	23.1	16.2 (15.9-16.5)	70.1 (68.9-71.3)	4.2 (3.9-4.4)	18.1 (17.0-19.2)	2.7 (2.5-2.9)	11.8 (10.8-12.7)
	65	1705	19.0	12.5 (12.2-12.8)	65.6 (64.1-67.0)	3.9 (3.6-4.1)	20.3 (19.0-21.7)	2.7 (2.5-2.9)	14.1 (12.9-15.3)
	70	1329	15.2	9.0 (8.7-9.3)	59.0 (57.1-60.9)	3.6 (3.3-3.9)	23.7 (21.9-25.4)	2.6 (2.4-2.9)	17.4 (15.8-18.9)
	75	1014	11.8	6.1 (5.8-6.4)	51.7 (49.2-54.2)	3.1 (2.9-3.4)	26.6 (24.3-28.9)	2.6 (2.3-2.8)	21.7 (19.5-23.9)
	80	619	8.7	3.8 (3.5-4.1)	43.7 (40.0-47.3)	2.6 (2.3-2.9)	29.8 (26.3-33.2)	2.3 (2.0-2.6)	26.6 (23.3-29.9)
	85+	243	6.2	2.4 (2.0-2.8)	39.0 (32.8-45.1)	2.0 (1.6-2.3)	31.7 (25.9-37.6)	1.8 (1.5-2.2)	29.3 (23.6-35.0)
2015	50	1920	32.9	24.8 (24.6-25.0)	75.2 (74.6-75.8)	5.3 (5.1-5.5)	16.1 (15.5-16.6)	2.9 (2.7-3.0)	8.7 (8.2-9.1)
	55	3035	28.4	20.7 (20.5-20.9)	72.7 (72.1-73.4)	4.9 (4.8-5.1)	17.3 (16.7-17.9)	2.8 (2.7-3.0)	9.9 (9.4-10.4)
	60	3360	24.2	16.9 (16.7-17.1)	69.8 (69.0-70.6)	4.6 (4.4-4.7)	18.8 (18.1-19.5)	2.8 (2.6-2.9)	11.4 (10.8-12.0)
	65	3224	20.2	13.2 (13.0-13.4)	65.4 (64.5-66.3)	4.2 (4.0-4.4)	20.9 (20.1-21.8)	2.8 (2.6-2.9)	13.7 (13.0-14.4)
	70	2535	16.3	9.7 (9.5-9.9)	59.5 (58.4-60.7)	3.8 (3.7-4.0)	23.5 (22.4-24.6)	2.8 (2.6-2.9)	17.0 (16.0-17.9)
	75	2051	12.6	6.6 (6.4-6.8)	52.2 (50.7-53.7)	3.3 (3.1-3.5)	26.0 (24.6-27.4)	2.8 (2.6-2.9)	21.8 (20.5-23.1)
	80	1294	9.4	3.9 (3.7-4.1)	41.8 (39.7-43.8)	2.8 (2.6-3.0)	30.0 (28.1-32.0)	2.6 (2.5-2.8)	28.2 (26.3-30.1)
	85+	972	6.6	2.3 (2.1-2.5)	34.6 (31.6-37.6)	2.1 (1.9-2.3)	31.5 (28.6-34.4)	2.3 (2.1-2.4)	33.9 (31.0-36.9)

^a Austria, Germany, France, Switzerland and Belgium.

Table A4: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Southern SHARE population^a in 2004-05 (Wave 1) and 2015 (Wave 6)

	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
2004-05	50	1189	32.3	20.3 (19.9-20.7)	62.9 (61.7-64.0)	7.3 (6.9-7.7)	22.6 (21.5-23.7)	4.7 (4.4-5.0)	14.5 (13.5-15.5)
	55	1218	27.8	16.5 (16.2-16.9)	59.5 (58.2-60.8)	6.7 (6.3-7.0)	23.9 (22.7-25.2)	4.6 (4.3-4.9)	16.6 (15.4-17.7)
	60	1148	23.5	13.0 (12.6-13.3)	55.3 (53.8-56.9)	6.0 (5.6-6.3)	25.5 (24.0-27.0)	4.5 (4.2-4.8)	19.1 (17.7-20.5)
	65	1031	19.3	9.6 (9.2-9.9)	49.5 (47.7-51.3)	5.4 (5.0-5.7)	27.9 (26.1-29.7)	4.4 (4.0-4.7)	22.6 (20.9-24.3)
	70	903	15.4	6.5 (6.1-6.8)	42.2 (40.0-44.4)	4.7 (4.3-5.0)	30.5 (28.3-32.8)	4.2 (3.9-4.5)	27.2 (25.0-29.5)
	75	575	11.8	4.2 (3.8-4.5)	35.3 (32.4-38.2)	3.8 (3.4-4.1)	32.1 (29.1-35.1)	3.8 (3.5-4.2)	32.6 (29.6-35.6)
	80	343	8.7	2.2 (1.9-2.6)	25.6 (21.6-29.5)	2.9 (2.5-3.2)	33.0 (28.7-37.3)	3.6 (3.2-4.0)	41.4 (36.9-45.9)
	85+	174	6.3	1.4 (1.0-1.8)	22.8 (16.6-29.0)	2.2 (1.7-2.6)	34.4 (27.4-41.5)	2.7 (2.2-3.2)	42.8 (35.4-50.1)
2015	50	1299	33.3	22.0 (21.7-22.2)	66.1 (65.3-66.8)	7.0 (6.7-7.2)	20.9 (20.2-21.6)	4.3 (4.1-4.5)	13.0 (12.4-13.6)
	55	2079	28.8	18.2 (18.0-18.5)	63.3 (62.5-64.2)	6.3 (6.0-6.5)	21.8 (21.0-22.6)	4.3 (4.1-4.5)	14.9 (14.2-15.6)
	60	2462	24.4	14.4 (14.1-14.6)	58.8 (57.8-59.8)	5.9 (5.6-6.1)	24.0 (23.0-24.9)	4.2 (4.0-4.4)	17.3 (16.4-18.1)
	65	2371	20.3	10.9 (10.7-11.2)	53.8 (52.7-54.9)	5.3 (5.1-5.5)	26.0 (24.9-27.1)	4.1 (3.9-4.3)	20.2 (19.2-21.2)
	70	1902	16.4	7.8 (7.6-8.1)	47.8 (46.4-49.2)	4.6 (4.4-4.8)	28.0 (26.7-29.4)	4.0 (3.8-4.2)	24.2 (22.9-25.4)
	75	1570	12.7	5.2 (5.0-5.4)	40.7 (38.9-42.4)	3.8 (3.6-4.0)	29.7 (28.0-31.4)	3.8 (3.6-4.0)	29.7 (27.9-31.4)
	80	1052	9.4	3.2 (2.9-3.4)	33.6 (31.2-35.9)	2.9 (2.7-3.1)	30.7 (28.3-33.1)	3.4 (3.1-3.6)	35.8 (33.3-38.3)
	85+	588	6.7	1.7 (1.5-2.0)	25.9 (22.3-29.4)	2.2 (1.9-2.5)	33.0 (29.2-36.8)	2.7 (2.5-3.0)	41.2 (37.2-45.1)

^aItaly, Spain and Greece.

Table A5: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Total SHARE population^a (2004-05 (Wave 1))

	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
Women	50	2180	34.2	21.7 (21.4-22.0)	63.3 (62.4-64.2)	7.8 (7.5-8.1)	22.9 (22.0-23.7)	4.7 (4.5-5.0)	13.9 (13.1-14.6)
	55	2195	29.7	17.8 (17.5-18.1)	59.9 (58.9-61.0)	7.2 (6.9-7.5)	24.2 (23.2-25.2)	4.7 (4.4-5.0)	15.9 (15.0-16.8)
	60	1942	25.2	14.1 (13.8-14.4)	55.9 (54.7-57.1)	6.5 (6.2-6.8)	25.7 (24.6-26.9)	4.6 (4.4-4.9)	18.4 (17.3-19.4)
	65	1744	20.8	10.6 (10.3-10.9)	50.9 (49.6-52.3)	5.7 (5.5-6.0)	27.5 (26.2-28.9)	4.5 (4.2-4.7)	21.5 (20.2-22.8)
	70	1419	16.6	7.3 (7.0-7.6)	43.9 (42.2-45.6)	5.1 (4.8-5.3)	30.4 (28.7-32.1)	4.3 (4.0-4.6)	25.7 (24.1-27.4)
	75	1097	12.7	4.8 (4.5-5.1)	37.6 (35.5-39.8)	4.1 (3.9-4.4)	32.5 (30.4-34.7)	3.8 (3.5-4.1)	29.8 (27.7-32.0)
	80	693	9.3	2.7 (2.5-3.0)	29.6 (26.7-32.5)	3.2 (2.9-3.4)	34.0 (30.9-37.1)	3.4 (3.1-3.7)	36.4 (33.3-39.5)
	85+	354	6.6	1.7 (1.4-2.0)	25.7 (21.1-30.2)	2.3 (2.0-2.6)	35.3 (30.3-40.3)	2.6 (2.2-2.9)	39.0 (33.9-44.1)
Men	50	1877	29.2	23.9 (23.7-24.2)	81.8 (81.0-82.6)	3.4 (3.2-3.6)	11.6 (10.9-12.3)	1.9 (1.8-2.1)	6.6 (6.0-7.1)
	55	1907	25.0	19.8 (19.6-20.1)	79.4 (78.5-80.3)	3.3 (3.1-3.5)	13.0 (12.2-13.8)	1.9 (1.7-2.1)	7.6 (6.9-8.2)
	60	1774	20.9	16.0 (15.8-16.3)	76.8 (75.7-78.0)	3.0 (2.8-3.2)	14.4 (13.4-15.3)	1.8 (1.7-2.0)	8.8 (8.0-9.6)
	65	1601	17.0	12.3 (12.1-12.6)	72.5 (71.1-73.9)	2.9 (2.7-3.1)	17.0 (15.8-18.2)	1.8 (1.6-2.0)	10.5 (9.5-11.6)
	70	1307	13.5	9.0 (8.7-9.2)	66.8 (64.9-68.6)	2.7 (2.5-2.9)	20.1 (18.5-21.7)	1.8 (1.6-2.0)	13.2 (11.8-14.6)
	75	903	10.3	6.2 (6.0-6.5)	60.5 (58.0-63.1)	2.3 (2.0-2.5)	21.9 (19.7-24.1)	1.8 (1.6-2.0)	17.5 (15.5-19.6)
	80	490	7.6	4.0 (3.7-4.3)	52.7 (48.8-56.6)	1.9 (1.6-2.2)	25.0 (21.6-28.4)	1.7 (1.4-1.9)	22.3 (19.0-25.5)
	85+	215	5.6	2.7 (2.4-3.1)	48.9 (42.3-55.6)	1.6 (1.2-1.9)	28.3 (22.3-34.3)	1.3 (1.0-1.6)	22.7 (17.1-28.3)

^a Denmark, Sweden, Austria, Germany, France, Switzerland, Belgium, Italy, Spain and Greece.

Table A6: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Total SHARE population^a (2015 (Wave 6))

Gender	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
Women	50	2339	35.4	22.6 (22.4-22.8)	63.8 (63.1-64.4)	7.9 (7.7-8.1)	22.3 (21.7-23.0)	4.9 (4.7-5.1)	13.9 (13.4-14.4)
	55	3458	30.8	18.7 (18.5-18.9)	60.8 (60.1-61.5)	7.2 (7.0-7.4)	23.5 (22.9-24.2)	4.8 (4.6-5.0)	15.7 (15.1-16.2)
	60	3785	26.3	15.0 (14.8-15.2)	56.9 (56.1-57.7)	6.7 (6.5-6.9)	25.4 (24.6-26.1)	4.7 (4.5-4.8)	17.7 (17.1-18.4)
	65	3669	21.9	11.4 (11.2-11.6)	51.9 (51.0-52.8)	6.0 (5.8-6.2)	27.3 (26.4-28.2)	4.6 (4.4-4.7)	20.8 (19.9-21.6)
	70	2890	17.7	8.1 (7.9-8.3)	45.8 (44.7-46.9)	5.2 (5.0-5.4)	29.2 (28.1-30.3)	4.4 (4.2-4.6)	25.0 (24.0-26.1)
	75	2279	13.7	5.3 (5.1-5.4)	38.3 (37.0-39.7)	4.2 (4.0-4.4)	30.9 (29.5-32.3)	4.2 (4.0-4.4)	30.8 (29.4-32.2)
	80	1530	10.1	3.0 (2.9-3.2)	30.3 (28.5-32.1)	3.3 (3.1-3.4)	32.3 (30.5-34.2)	3.8 (3.6-4.0)	37.4 (35.5-39.3)
	85+	1094	7.1	1.7 (1.5-1.9)	23.9 (21.4-26.4)	2.3 (2.1-2.5)	33.1 (30.3-35.9)	3.0 (2.8-3.2)	43.0 (40.0-45.9)
Men	50	1557	31.0	25.4 (25.2-25.6)	82.0 (81.5-82.6)	3.8 (3.6-3.9)	12.2 (11.7-12.7)	1.8 (1.7-1.9)	5.7 (5.4-6.1)
	55	2632	26.6	21.3 (21.1-21.5)	80.2 (79.6-80.8)	3.5 (3.3-3.6)	13.1 (12.6-13.6)	1.8 (1.7-1.9)	6.7 (6.3-7.1)
	60	3261	22.4	17.4 (17.2-17.6)	77.7 (76.9-78.4)	3.2 (3.1-3.4)	14.5 (13.8-15.1)	1.8 (1.7-1.9)	7.9 (7.4-8.4)
	65	3360	18.5	13.7 (13.5-13.9)	74.0 (73.1-74.9)	3.0 (2.9-3.2)	16.5 (15.7-17.2)	1.8 (1.6-1.9)	9.5 (8.9-10.2)
	70	2695	14.8	10.3 (10.1-10.5)	69.5 (68.4-70.6)	2.8 (2.7-3.0)	19.0 (18.1-20.0)	1.7 (1.6-1.8)	11.5 (10.7-12.3)
	75	2151	11.4	7.2 (7.1-7.4)	63.5 (62.1-65.0)	2.5 (2.3-2.6)	21.7 (20.4-23.0)	1.7 (1.6-1.8)	14.8 (13.7-15.9)
	80	1316	8.4	4.6 (4.4-4.8)	55.0 (52.9-57.2)	2.2 (2.0-2.3)	25.9 (24.0-27.8)	1.6 (1.5-1.7)	19.1 (17.4-20.8)
	85+	843	6.0	2.9 (2.7-3.1)	47.8 (44.4-51.1)	1.7 (1.6-1.9)	29.2 (26.1-32.2)	1.4 (1.2-1.6)	23.1 (20.2-25.9)

^a Denmark, Sweden, Austria, Germany, France, Switzerland, Belgium, Italy, Spain and Greece.

Table A7: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Northern SHARE population^a (2004-05 (Wave 1))

Gender	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
Women	50	411	33.1	24.4 (23.8-25.0)	73.6 (71.8-75.5)	6.2 (5.7-6.8)	18.8 (17.1-20.6)	2.5 (2.1-2.9)	7.5 (6.2-8.8)
	55	479	28.6	20.3 (19.7-20.9)	70.9 (68.8-73.0)	5.9 (5.3-6.4)	20.5 (18.5-22.5)	2.5 (2.0-2.9)	8.6 (7.1-10.1)
	60	382	24.2	16.5 (15.9-17.1)	68.1 (65.7-70.6)	5.4 (4.8-5.9)	22.1 (19.7-24.5)	2.4 (1.9-2.8)	9.7 (7.9-11.5)
	65	307	20.0	12.8 (12.2-13.4)	63.8 (60.9-66.8)	4.9 (4.3-5.5)	24.6 (21.7-27.4)	2.3 (1.9-2.8)	11.6 (9.4-13.8)
	70	250	16.1	9.2 (8.7-9.8)	57.5 (53.8-61.2)	4.6 (4.0-5.2)	28.5 (24.9-32.1)	2.2 (1.8-2.7)	14.0 (11.1-16.8)
	75	213	12.5	6.2 (5.6-6.8)	49.5 (44.8-54.3)	4.0 (3.5-4.6)	32.4 (27.8-37.1)	2.2 (1.8-2.7)	18.0 (14.1-21.9)
	80	116	9.2	3.6 (3.0-4.2)	39.0 (32.4-45.6)	3.4 (2.8-4.0)	37.2 (30.6-43.8)	2.2 (1.7-2.7)	23.8 (17.9-29.7)
	85	93	6.6	2.2 (1.6-2.8)	33.8 (24.2-43.5)	2.7 (2.1-3.4)	41.6 (31.5-51.6)	1.6 (1.0-2.2)	24.6 (15.8-33.4)
Men	50	364	29.3	26.2 (25.7-26.6)	89.2 (87.8-90.6)	2.1 (1.7-2.4)	7.0 (5.8-8.2)	1.1 (0.8-1.4)	3.8 (2.9-4.7)
	55	391	24.9	21.8 (21.4-22.2)	87.3 (85.6-89.0)	2.1 (1.7-2.4)	8.3 (6.9-9.7)	1.1 (0.8-1.4)	4.4 (3.3-5.5)
	60	368	20.7	17.7 (17.3-18.1)	85.4 (83.3-87.4)	2.0 (1.6-2.3)	9.4 (7.7-11.2)	1.1 (0.8-1.4)	5.2 (3.9-6.5)
	65	302	16.8	13.9 (13.5-14.3)	82.5 (79.9-85.0)	1.9 (1.5-2.3)	11.2 (9.1-13.4)	1.1 (0.8-1.3)	6.3 (4.6-8.0)
	70	244	13.3	10.4 (9.9-10.8)	78.2 (74.8-81.7)	1.8 (1.4-2.2)	13.7 (10.8-16.6)	1.1 (0.8-1.4)	8.1 (5.7-10.4)
	75	198	10.1	7.4 (7.0-7.9)	73.7 (69.0-78.4)	1.7 (1.3-2.1)	16.5 (12.5-20.6)	1.0 (0.7-1.3)	9.8 (6.5-13.0)
	80	105	7.4	5.2 (4.6-5.7)	70.2 (63.1-77.3)	1.3 (0.9-1.8)	18.1 (12.1-24.1)	0.9 (0.5-1.2)	11.7 (6.7-16.8)
	85	59	5.3	3.5 (2.8-4.1)	65.7 (53.6-77.8)	1.1 (0.6-1.7)	21.7 (11.2-32.2)	0.7 (0.2-1.1)	12.7 (4.2-21.1)

^a Denmark and Sweden.

Table A8: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Northern SHARE population^a (2015 (Wave 6))

Gender	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
Women	50	392	34.6	26.5 (26.0-27.0)	76.6 (75.3-78.0)	5.5 (5.0-5.9)	15.8 (14.5-17.0)	2.6 (2.3-2.9)	7.6 (6.6-8.5)
	55	543	29.9	22.2 (21.7-22.6)	74.2 (72.7-75.7)	5.1 (4.7-5.5)	17.1 (15.8-18.5)	2.6 (2.3-2.9)	8.7 (7.6-9.8)
	60	651	25.4	18.1 (17.6-18.5)	71.1 (69.4-72.9)	4.8 (4.4-5.2)	18.9 (17.3-20.5)	2.5 (2.2-2.9)	10.0 (8.7-11.3)
	65	747	21.1	14.2 (13.8-14.7)	67.4 (65.4-69.5)	4.3 (3.9-4.7)	20.6 (18.7-22.4)	2.5 (2.2-2.9)	12.0 (10.4-13.6)
	70	602	17.0	10.6 (10.1-11.0)	62.2 (59.7-64.7)	4.0 (3.6-4.4)	23.3 (21.0-25.7)	2.5 (2.1-2.8)	14.5 (12.5-16.5)
	75	398	13.2	7.3 (6.8-7.7)	55.2 (51.9-58.5)	3.5 (3.1-3.9)	26.8 (23.8-29.9)	2.4 (2.0-2.7)	18.0 (15.3-20.7)
	80	270	9.7	4.6 (4.1-5.0)	46.9 (42.4-51.4)	2.8 (2.4-3.2)	29.2 (25.0-33.4)	2.3 (1.9-2.7)	23.9 (19.9-27.9)
	85	206	6.9	2.6 (2.2-3.1)	38.2 (31.6-44.9)	2.2 (1.8-2.7)	32.3 (25.9-38.7)	2.0 (1.6-2.4)	29.5 (23.3-35.7)
Men	50	285	31.3	27.9 (27.6-28.3)	89.1 (88.1-90.2)	2.1 (1.8-2.4)	6.7 (5.8-7.6)	1.3 (1.1-1.5)	4.2 (3.5-4.9)
	55	433	26.9	23.6 (23.2-23.9)	87.7 (86.5-88.9)	2.0 (1.7-2.3)	7.4 (6.5-8.4)	1.3 (1.1-1.5)	4.9 (4.0-5.7)
	60	573	22.5	19.3 (19.0-19.6)	85.7 (84.3-87.1)	1.9 (1.7-2.2)	8.6 (7.4-9.7)	1.3 (1.1-1.5)	5.7 (4.7-6.7)
	65	687	18.5	15.4 (15.1-15.7)	83.2 (81.5-84.9)	1.8 (1.5-2.1)	9.7 (8.3-11.1) ¹	1.3 (1.1-1.5)	7.1 (5.8-8.3)
	70	546	14.6	11.7 (11.4-12.0)	79.9 (77.7-82.2)	1.6 (1.4-1.9)	11.1 (9.3-12.8)	1.3 (1.1-1.6)	9.0 (7.4-10.6)
	75	411	11.2	8.4 (8.1-8.7)	75.0 (71.9-78.0)	1.5 (1.2-1.7)	12.9 (10.5-15.4)	1.4 (1.1-1.6)	12.1 (9.7-14.4)
	80	230	8.1	5.7 (5.3-6.0)	69.7 (65.2-74.2)	1.3 (1.0-1.5)	15.4 (11.9-19.0)	1.2 (0.9-1.5)	14.9 (11.4-18.4)
	85	171	5.7	3.5 (3.1-3.9)	61.6 (54.3-68.9)	1.0 (0.7-1.3)	17.4 (11.7-23.1)	1.2 (0.8-1.5)	21.0 (14.9-27.1)

^a Denmark and Sweden.

Table A9: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Central SHARE population^a (2004-05 (Wave 1))

Gender	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
Women	50	1092	34.0	23.8 (23.4-24.2)	69.9 (68.7-71.2)	6.5 (6.1-6.9)	19.2 (18.1-20.4)	3.7 (3.3-4.0)	10.8 (9.8-11.8)
	55	1069	29.5	19.7 (19.3-20.1)	66.8 (65.4-68.2)	6.1 (5.7-6.5)	20.6 (19.3-22.0)	3.7 (3.4-4.1)	12.6 (11.4-13.8)
	60	945	25.0	15.8 (15.3-16.2)	62.9 (61.3-64.6)	5.6 (5.2-6.0)	22.2 (20.7-23.8)	3.7 (3.4-4.1)	14.8 (13.4-16.2)
	65	887	20.7	12.1 (11.7-12.5)	58.4 (56.4-60.3)	5.0 (4.6-5.4)	24.1 (22.2-26.0)	3.6 (3.3-4.0)	17.6 (15.8-19.3)
	70	708	16.5	8.5 (8.1-8.9)	51.5 (49.0-54.0)	4.5 (4.1-4.9)	27.2 (24.9-29.6)	3.5 (3.2-3.9)	21.3 (19.0-23.5)
	75	562	12.7	5.6 (5.2-6.0)	44.3 (41.1-47.5)	3.8 (3.4-4.2)	30.2 (27.1-33.3)	3.2 (2.9-3.6)	25.5 (22.5-28.5)
	80	373	9.2	3.3 (2.9-3.7)	35.7 (31.3-40.2)	3.0 (2.6-3.4)	32.8 (28.4-37.3)	2.9 (2.5-3.3)	31.4 (27.0-35.9)
	85	157	6.5	2.0 (1.5-2.5)	31.0 (23.7-38.2)	2.2 (1.7-2.6)	33.4 (26.0-40.8)	2.3 (1.8-2.8)	35.6 (28.1-43.1)
Men	50	1001	29.0	24.6 (24.3-24.9)	85.0 (83.9-86.1)	2.8 (2.5-3.1)	9.6 (8.7-10.6)	1.6 (1.4-1.8)	5.4 (4.7-6.2)
	55	945	24.8	20.5 (20.2-20.9)	82.9 (81.6-84.2)	2.7 (2.4-3.0)	10.9 (9.8-12.0)	1.5 (1.3-1.8)	6.2 (5.3-7.1)
	60	873	20.8	16.7 (16.4-17.1)	80.6 (79.1-82.2)	2.5 (2.3-2.8)	12.2 (10.9-13.6)	1.5 (1.3-1.7)	7.1 (6.0-8.2)
	65	818	16.9	13.0 (12.7-13.4)	76.9 (74.9-78.9)	2.5 (2.2-2.8)	14.6 (12.9-16.4)	1.4 (1.2-1.7)	8.4 (7.1-9.8)
	70	621	13.4	9.7 (9.3-10.0)	71.9 (69.2-74.5)	2.4 (2.1-2.7)	17.6 (15.3-19.9)	1.4 (1.2-1.7)	10.5 (8.6-12.3)
	75	452	10.3	6.8 (6.4-7.2)	65.9 (62.1-69.6)	2.0 (1.7-2.4)	19.9 (16.6-23.1)	1.5 (1.2-1.7)	14.2 (11.5-17.0)
	80	246	7.6	4.7 (4.3-5.1)	61.6 (55.8-67.3)	1.7 (1.3-2.1)	22.7 (17.6-27.7)	1.2 (0.9-1.5)	15.8 (11.6-20.0)
	85	86	5.6	3.3 (2.7-3.9)	58.6 (48.2-69.0)	1.6 (1.0-2.1)	27.6 (18.2-37.1)	0.8 (0.4-1.2)	13.8 (6.5-21.1)

^a Austria, Germany, France, Switzerland and Belgium.

Table A10: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Central SHARE population^a (2015 (Wave 6))

Gender	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
Women	50	1150	35.1	24.0 (23.7-24.4)	68.5 (67.6-69.3)	7.0 (6.7-7.3)	19.9 (19.1-20.7)	4.1 (3.8-4.3)	11.6 (11.0-12.3)
	55	1690	30.5	20.0 (19.7-20.3)	65.7 (64.7-66.6)	6.5 (6.2-6.8)	21.3 (20.3-22.2)	4.0 (3.8-4.2)	13.1 (12.3-13.8)
	60	1823	26.1	16.2 (16.0-16.5)	62.3 (61.2-63.4)	6.0 (5.7-6.3)	22.9 (21.9-24.0)	3.8 (3.6-4.1)	14.7 (13.8-15.6)
	65	1692	21.7	12.5 (12.3-12.8)	57.7 (56.4-59.0)	5.4 (5.1-5.7)	24.8 (23.6-26.1)	3.8 (3.6-4.0)	17.5 (16.4-18.6)
	70	1320	17.6	9.1 (8.8-9.3)	51.5 (49.9-53.1)	4.7 (4.5-5.0)	26.9 (25.4-28.4)	3.8 (3.6-4.1)	21.6 (20.2-23.0)
	75	1097	13.6	6.0 (5.7-6.2)	43.8 (41.8-45.7)	3.9 (3.7-4.2)	28.9 (27.0-30.8)	3.7 (3.5-4.0)	27.3 (25.4-29.2)
	80	706	10.0	3.4 (3.2-3.7)	34.4 (31.9-37.0)	3.1 (2.9-3.4)	31.4 (28.8-34.0)	3.4 (3.2-3.7)	34.2 (31.5-36.8)
	85	581	7.0	1.9 (1.7-2.2)	27.5 (23.9-31.1)	2.2 (2.0-2.5)	31.9 (28.1-35.6)	2.8 (2.6-3.1)	40.7 (36.7-44.7)
Men	50	770	30.6	25.5 (25.3-25.8)	83.5 (82.7-84.2)	3.5 (3.3-3.8)	11.6 (10.9-12.3)	1.5 (1.4-1.7)	5.0 (4.5-5.4)
	55	1345	26.2	21.4 (21.1-21.6)	81.5 (80.6-82.4)	3.3 (3.1-3.5)	12.6 (11.9-13.4)	1.5 (1.4-1.7)	5.9 (5.3-6.4)
	60	1537	22.1	17.6 (17.4-17.8)	79.5 (78.5-80.5)	3.0 (2.8-3.2)	13.7 (12.7-14.6)	1.5 (1.4-1.7)	6.9 (6.2-7.6)
	65	1532	18.3	13.9 (13.7-14.1)	75.8 (74.5-77.0)	2.9 (2.7-3.1)	15.9 (14.7-17.0)	1.5 (1.4-1.7)	8.4 (7.5-9.2)
	70	1215	14.7	10.5 (10.3-10.7)	71.2 (69.6-72.8)	2.8 (2.6-3.0)	18.8 (17.4-20.2)	1.5 (1.3-1.6)	10.0 (8.9-11.1)
	75	954	11.3	7.4 (7.2-7.7)	65.4 (63.3-67.6)	2.4 (2.2-2.7)	21.6 (19.7-23.5)	1.5 (1.3-1.7)	13.0 (11.4-14.6)
	80	588	8.4	4.6 (4.3-4.9)	55.2 (52.0-58.3)	2.3 (2.1-2.5)	27.5 (24.7-30.4)	1.4 (1.2-1.7)	17.3 (14.9-19.8)
	85	391	6.0	2.9 (2.6-3.2)	48.5 (43.6-53.5)	1.8 (1.6-2.1)	30.7 (26.2-35.3)	1.2 (1.0-1.5)	20.7 (16.7-24.7)

^a Austria, Germany, France, Switzerland and Belgium.

Table A11: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Southern SHARE population^a (2004-05 (Wave 1))

Gender	Age at start of the interval	N	Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
Women	50	677	34.7	18.0 (17.4-18.5)	51.7 (50.1-53.3)	10.0 (9.4-10.6)	28.8 (27.1-30.5)	6.8 (6.2-7.3)	19.5 (17.9-21.0)
	55	647	30.1	14.5 (14.0-15.0)	48.2 (46.4-50.0)	9.0 (8.4-9.5)	29.8 (27.9-31.7)	6.6 (6.1-7.1)	22.0 (20.2-23.8)
	60	615	25.5	11.1 (10.6-11.6)	43.6 (41.6-45.6)	8.0 (7.4-8.6)	31.3 (29.1-33.5)	6.4 (5.9-6.9)	25.1 (23.0-27.1)
	65	550	21.1	8.0 (7.5-8.5)	38.0 (35.7-40.4)	6.9 (6.4-7.5)	32.9 (30.3-35.5)	6.1 (5.6-6.7)	29.0 (26.5-31.6)
	70	461	16.8	5.2 (4.7-5.7)	30.8 (28.1-33.6)	5.9 (5.4-6.4)	35.0 (31.8-38.2)	5.8 (5.2-6.3)	34.2 (31.0-37.4)
	75	322	12.9	3.3 (2.8-3.7)	25.3 (21.9-28.7)	4.6 (4.1-5.2)	36.0 (31.9-40.2)	5.0 (4.4-5.5)	38.7 (34.5-42.9)
	80	204	9.4	1.7 (1.3-2.1)	17.9 (13.5-22.4)	3.3 (2.8-3.9)	35.2 (29.4-41.0)	4.4 (3.8-5.0)	46.9 (40.9-52.9)
	85	104	6.7	1.0 (0.5-1.4)	14.3 (7.5-21.0)	2.5 (1.9-3.1)	37.2 (27.9-46.5)	3.2 (2.6-3.9)	48.5 (38.9-58.1)
Men	50	512	29.7	22.7 (22.2-23.1)	76.4 (74.9-77.9)	4.5 (4.1-4.9)	15.1 (13.8-16.5)	2.5 (2.2-2.8)	8.5 (7.3-9.6)
	55	571	25.3	18.6 (18.1-19.0)	73.5 (71.8-75.3)	4.2 (3.8-4.6)	16.7 (15.1-18.3)	2.5 (2.1-2.8)	9.8 (8.4-11.1)
	60	533	21.1	14.9 (14.5-15.3)	70.5 (68.5-72.6)	3.8 (3.4-4.2)	18.1 (16.2-20.0)	2.4 (2.1-2.7)	11.4 (9.7-13.0)
	65	481	17.2	11.2 (10.8-11.6)	65.2 (62.6-67.8)	3.6 (3.2-4.0)	21.1 (18.7-23.4)	2.4 (2.0-2.7)	13.7 (11.7-15.8)
	70	442	13.5	8.0 (7.5-8.4)	58.8 (55.5-62.1)	3.3 (2.8-3.7)	24.1 (21.0-27.2)	2.3 (1.9-2.7)	17.1 (14.3-19.9)
	75	253	10.3	5.3 (4.8-5.8)	51.5 (46.9-56.1)	2.7 (2.2-3.1)	25.8 (21.6-30.0)	2.3 (1.9-2.8)	22.7 (18.6-26.7)
	80	139	7.6	2.9 (2.4-3.4)	38.6 (31.8-45.4)	2.2 (1.7-2.7)	29.4 (23.0-35.8)	2.4 (1.9-2.9)	32.0 (25.4-38.5)
	85	70	5.6	2.0 (1.4-2.6)	35.6 (24.4-46.8)	1.7 (1.1-2.3)	30.2 (19.4-40.9)	1.9 (1.3-2.5)	34.2 (23.1-45.3)

^a Italy, Spain and Greece.

Table A12: Life expectancy (LE) by frailty state in women and men by 5-year age group in the Southern SHARE population^a (2015 (Wave 6))

Gender	Age at start of the interval	N	Total Life Expectancy	Non-Frail		Pre-Frail		Frail	
				LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy	LE (95% CI)	% of total life expectancy
Women	50	797	36.0	19.5 (19.2-19.9)	54.3 (53.3-55.4)	9.8 (9.4-10.) ²	27.2 (26.0-28.3)	6.6 (6.3-7.0)	18.5 (17.5-19.5)
	55	1225	31.3	15.9 (15.6-16.3)	50.9 (49.8-52.1)	8.8 (8.4-9.2)	28.1 (26.9-29.4)	6.6 (6.2-6.9)	20.9 (19.8-22.1)
	60	1311	26.7	12.3 (11.9-12.6)	46.0 (44.7-47.3)	8.1 (7.7-8.5)	30.2 (28.8-31.6)	6.4 (6.0-6.7)	23.8 (22.5-25.1)
	65	1230	22.3	9.0 (8.7-9.3)	40.4 (38.9-41.9)	7.2 (6.8-7.6)	32.2 (30.6-33.9)	6.1 (5.7-6.5)	27.4 (25.8-29.0)
	70	968	18.0	6.1 (5.8-6.4)	33.9 (32.1-35.6)	6.1 (5.7-6.5)	33.9 (31.9-36.0)	5.8 (5.4-6.1)	32.2 (30.2-34.2)
	75	784	13.9	3.7 (3.4-4.0)	26.8 (24.7-29.0)	4.8 (4.5-5.2)	34.7 (32.2-37.2)	5.3 (5.0-5.7)	38.5 (35.9-41.1)
	80	554	10.2	2.1 (1.8-2.4)	20.4 (17.7-23.2)	3.5 (3.2-3.9)	34.6 (31.1-38.0)	4.6 (4.2-5.0)	45.0 (41.4-48.6)
	85	307	7.2	1.0 (0.7-1.3)	14.3 (10.3-18.2)	2.6 (2.2-3.0)	35.9 (30.5-41.3)	3.6 (3.2-4.0)	49.9 (44.3-55.4)
Men	50	502	31.5	24.9 (24.5-25.2)	78.9 (77.9-79.9)	4.4 (4.1-4.7)	13.9 (13.0-14.8)	2.3 (2.1-2.5)	7.2 (6.5-7.9)
	55	854	27.0	20.9 (20.6-21.1)	77.1 (76.0-78.2)	4.0 (3.7-4.2)	14.6 (13.7-15.6)	2.2 (2.0-2.4)	8.3 (7.5-9.0)
	60	1151	22.8	16.8 (16.5-17.1)	73.6 (72.3-74.9)	3.8 (3.5-4.0)	16.6 (15.5-17.7)	2.2 (2.0-2.4)	9.8 (8.9-10.) ⁷
	65	1141	18.8	13.1 (12.9-13.4)	70.0 (68.5-71.5)	3.4 (3.2-3.7)	18.3 (17.0-19.7)	2.2 (2.0-2.4)	11.7 (10.5-12.8)
	70	934	15.0	9.8 (9.5-10.1)	65.3 (63.4-67.2)	3.1 (2.8-3.3)	20.6 (18.9-22.2)	2.1 (1.9-2.3)	14.1 (12.6-15.6)
	75	786	11.5	6.8 (6.5-7.1)	59.1 (56.5-61.6)	2.6 (2.4-2.9)	23.0 (20.8-25.2)	2.1 (1.8-2.3)	17.9 (15.9-20.0)
	80	498	8.5	4.5 (4.2-4.8)	52.8 (49.2-56.4)	2.1 (1.8-2.4)	24.9 (21.7-28.0)	1.9 (1.6-2.2)	22.3 (19.2-25.4)
	85	281	6.0	2.7 (2.3-3.0)	44.3 (38.5-50.2)	1.7 (1.4-2.0)	28.3 (23.1-33.6)	1.7 (1.3-2.0)	27.3 (22.1-32.5)

^a Italy, Spain and Greece.