

Study of perinatal transmission of SARS-CoV-2 in a Mexican public hospital

Adriana Ramírez-Rosas, Tizziani Benitez-Guerrero, Karina Corona-Cervantes, Juan Manuel Vélez-Ixta, Norma Gabriela Zavala-Torres, Jazmin Cuenca-Leija, Sarahi Martínez- Pichardo, María Esther Landero-Montes-de-Oca, Fernando Guadalupe Bastida-González, Paola Berenice Zárate-Segura, Jaime García-Mena.

Supplementary material

Table S1

Detection of SARS-CoV-2 RNA by qPCR in nasopharyngeal swabs and human milk samples.

Sample	N	Positive	Negative
mother nasopharyngeal swab	133	16 (12.03%)	117 (87.97%)
colostrum/human milk sample	140	3 (2.14%)	137 (97.86%)
neonate nasopharyngeal swab	131 ^a	11 (8.40%)	120 (91.60%)

Table S2

Detection of SARS-CoV-2 RNA by ddPCR in nasopharyngeal swabs and human milk samples.

Sample	N	Positive	negative
mother nasopharyngeal swab	133	41 (30.83%)	92 (69.17%)
colostrum/human milk sample	140	20 (14.29%)	120 (85.71%)
neonate nasopharyngeal swab	131	24 (18.32%)	107 (81.68%)