
Dellepiane et al, Kidney Medicine, "Acute Kidney Injury in Patients Hospitalized With COVID-19 in New York City: Temporal Trends From March 2020 to April 2021"

Figure S1. Acute kidney injury stages over time according to the kidney disease improving global outcomes (KDIGO) classification.



Dellepiane et al, Kidney Medicine, "Acute Kidney Injury in Patients Hospitalized With COVID-19 in New York City: Temporal Trends From March 2020 to April 2021"

Month	Imputed creatinine (N(%))
March 2020	1152 (60.73)
April 2020	2065 (63.56)
May 2020	301 (55.13)
June 2020	74 (42.29)
July 2020	55 (43.65)
August 2020	72 (56.69)
September 2020	104 (53.33)
October 2020	187 (67.75)
November 2020	234 (62.9)
December 2020	571 (60.62)
January 2021	709 (59.48)
February 2021	657 (60.44)
March 2021	599 (60.02)
April 2021	390 (63)

Table S1. Percentage of imputed creatinine values over time

Dellepiane et al, Kidney Medicine, "Acute Kidney Injury in Patients Hospitalized With COVID-19 in New York City: Temporal Trends From March 2020 to April 2021"

Variable	Value
Age (mean, std)	70.08 (15.29)
eGFR from baseline Cr. (mean, std)	67.65 (36.83)
CKD (n, %)	303 (5.5%)
Gender (n, %)	
Male	3156 (57.26%)
Female	2356 (42.74%)
Race (n, %)	
Other	1913 (34.80%)
White	1518 (27.62%)
Black	1492 (27.14%)
Asian	305 (5.55%)
Unknown	264 (4.80%)

Table S2. Demographic characteristics of studied individuals

Legend:

CKD: chronic kidney disease, cr: creatinine, eGFR: estimated glomerular filtration rate, Std: standard deviation