

## Supplementary Figures

# Multi-ancestry genome-wide gene-sleep interactions identify novel loci for blood pressure



Wang H, et al.

## Contents

Supplementary Fig. 1. Miami plots of genome-wide gene-sleep interaction analyses in stage 1 MULTI ancestry group.....	2
Supplementary Fig. 2. QQ plots of genome-wide gene-sleep interaction analyses in stage 1 MULTI ancestry group.....	3
Supplementary Fig. 3. Miami plots of genome-wide gene-sleep interaction analyses in stage 1 EUR group.....	4
Supplementary Fig. 4. QQ plots of genome-wide gene-sleep interaction analyses in stage 1 EUR group..	5
Supplementary Fig. 5. Miami plots of genome-wide gene-sleep interaction analyses in stage 1 AFR group.....	6
Supplementary Fig. 6. QQ plots of genome-wide gene-sleep interaction analyses in stage 1 AFR group. .	7
Supplementary Fig. 7. Regional association plots of $P_{\text{joint}}$ and $P_{\text{int}}$ for 3 replicated novel loci in stage 1+2 MULTI ancestry analyses (Table 1). .....	8
Supplementary Fig. 8. Regional association plots of $P_{\text{joint}}$ and $P_{\text{int}}$ for 3 novel loci in stage 1 AFR analyses (Supplementary Table 8).....	9
Supplementary Fig. 9. Miami plots of genome-wide gene-sleep interaction analyses in stage 1+2 MULTI ancestry group.....	10
Supplementary Fig. 10. QQ plots of genome-wide gene-sleep interaction analyses in stage 1+2 MULTI ancestry group.....	11
Supplementary Fig. 11. Miami plots of genome-wide gene-sleep interaction analyses in stage 1+2 EUR group.....	12
Supplementary Fig. 12. QQ plots of genome-wide gene-sleep interaction analyses in stage 1+2 EUR group.....	13
Supplementary Fig. 13. Regional association plots of $P_{\text{joint}}$ and $P_{\text{int}}$ for 20 novel loci in stage 1+2 analyses (Supplementary Table 10). .....	14


## Supplementary Fig. 1. Miami plots of genome-wide gene-sleep interaction analyses in stage 1 MULTI ancestry group.

Upper panel are results without adjusting for BMI, lower panel are results adjusted for BMI. The minimum p-values across four BP traits were used to generate each plot. The orange points correspond to significant loci ( $P < 5 \times 10^{-8}$ ) overlapped with previously reported BP loci (distance  $< 1\text{Mb}$ ), and the red points correspond to novel significant loci.




Supplementary Fig. 2. QQ plots of genome-wide gene-sleep interaction analyses in stage 1 MULTI ancestry group.


LTST-SBP




LTST-DBP




LTST-MAP




LTST-PP




STST-SBP




STST-DBP



STST-MAP




STST-PP




### Supplementary Fig. 3. Miami plots of genome-wide gene-sleep interaction analyses in stage 1 EUR group.

Upper panel are results without adjusting for BMI, lower panel are results adjusted for BMI. The minimum p-values across four BP traits were used to generate each plot. The orange points correspond to significant loci ( $P < 5 \times 10^{-8}$ ) overlapped with previously reported BP loci (distance  $< 1\text{Mb}$ ), and the red points correspond to novel significant loci.




Supplementary Fig. 4. QQ plots of genome-wide gene-sleep interaction analyses in stage 1 EUR group.


LTST-SBP




LTST-DBP




LTST-MAP




LTST-PP




STST-SBP




STST-DBP



STST-MAP




STST-PP




## Supplementary Fig. 5. Miami plots of genome-wide gene-sleep interaction analyses in stage 1 AFR group.

Upper panel are results without adjusting for BMI, lower panel are results adjusted for BMI. The minimum p-values across four BP traits were used to generate each plot. The orange points correspond to significant loci ( $P < 5 \times 10^{-8}$ ) overlapped with previously reported BP loci (distance  $< 1\text{Mb}$ ), and the red points correspond to novel significant loci.




Supplementary Fig. 6. QQ plots of genome-wide gene-sleep interaction analyses in stage 1 AFR group.


LTST-SBP




LTST-DBP




LTST-MAP




LTST-PP




STST-SBP




STST-DBP



STST-MAP





STST-PP





Supplementary Fig. 7. Regional association plots of  $P_{\text{joint}}$  and  $P_{\text{int}}$  for 3 replicated novel loci in stage 1+2 MULTI ancestry analyses (Table 1).



Locus 1. LTST- MAP



Locus 2. STST- DBP





Locus 3. STST- PP





**Supplementary Fig. 8. Regional association plots of  $P_{\text{joint}}$  and  $P_{\text{int}}$  for 3 novel loci in stage 1 AFR analyses (Supplementary Table 8).**



**Locus 1. LTST- PP**



**Locus 2. LTST-SBP**




**Locus 3. LTST-PP**




### Supplementary Fig. 9. Miami plots of genome-wide gene-sleep interaction analyses in stage 1+2 MULTI ancestry group.

Upper panel are results without adjusting for BMI, lower panel are results adjusted for BMI. The minimum p-values across four BP traits were used to generate each plot. The orange points correspond to significant loci ( $P < 5 \times 10^{-8}$ ) overlapped with previously reported BP loci (distance  $< 1\text{Mb}$ ), and the red points correspond to novel significant loci.




Supplementary Fig. 10. QQ plots of genome-wide gene-sleep interaction analyses in stage 1+2 MULTI ancestry group.


LTST-SBP




LTST-DBP




LTST-MAP




LTST-PP




STST-SBP




STST-DBP



STST-MAP




STST-PP




## Supplementary Fig. 11. Miami plots of genome-wide gene-sleep interaction analyses in stage 1+2 EUR group.

Upper panel are results without adjusting for BMI, lower panel are results adjusted for BMI. The minimum p-values across four BP traits were used to generate each plot. The orange points correspond to significant loci ( $P < 5 \times 10^{-8}$ ) overlapped with previously reported BP loci (distance  $< 1\text{Mb}$ ), and the red points correspond to novel significant loci.




Supplementary Fig. 12. QQ plots of genome-wide gene-sleep interaction analyses in stage 1+2 EUR group.


LTST-SBP




LTST-DBP




LTST-MAP




LTST-PP




STST-SBP




STST-DBP



STST-MAP





STST-PP





Supplementary Fig. 13. Regional association plots of  $P_{joint}$  and  $P_{int}$  for 20 novel loci in stage 1+2 analyses (Supplementary Table 10).



Locus 1. LTST- PP





Locus 2. LTST- SBP





Locus 3. LTST- MAP





#### Locus 4. LTST- PP





#### Locus 5. LTST- SBP





#### Locus 6. LTST- SBP





### Locus 7. LTST- PP




### Locus 8. LTST- DBP




### Locus 9. LTST- PP




### Locus 10. STST- SBP





### Locus 11. STST- PP





### Locus 12. STST- MAP





### Locus 13. STST- PP




### Locus 14. STST- SBP




### Locus 15. STST- DBP




### Locus 16. STST- MAP




### Locus 17. STST- PP



### Locus 18. STST- DBP



### Locus 19. STST- SBP



### Locus 20. STST- PP

