

Supplementary Table S1. Baseline characteristics of included and excluded patients

Variable	Included, n=74	Excluded, n=26	<i>p</i>
Age, mean ± SD	81.5 (5.8)	79.9 (6.4)	0.35
Male, n (%)	28 (37.8)	14 (53.8)	0.16
Living at home, n (%)	65 (87.8)	21 (80.8)	0.65
Data missing	0	3	
No or only primary education, n (%)	29 (39.2)	10 (38.5)	0.98
Data missing	2	1	
Number of (i)ADL dependencies, median (IQR)	2.0 (1.0-4.25)	1.5 (0.0-5.0)	0.95
Subjective memory complaints, n (%)	24 (32.4)	9 (34.6)	0.87
Data missing	1	0	
Dementia, n (%)	0 (0)	1 (3.8)	0.09
Prior delirium, n (%)	12 (16.2)	5 (19.2)	0.73
MMSE, median (IQR)	28.0 (27.0-29.0)	27.0 (24.3-28.5)	0.04
Data missing	0	1	
Charlson Comorbidity Index, median (IQR)	2.0 (1.0-3.0)	2.0 (1.0-3.0)	0.84
Active cancer, n (%)	7 (9.5)	4 (15.4)	0.41
Coronary artery disease, n (%)	29 (39.2)	16 (61.5)	0.05
Chronic obstructive pulmonary disease, n (%)	24 (32.4)	9 (34.6)	0.84
Diabetes mellitus, n (%)	16 (21.6)	9 (34.6)	0.19
Atrial fibrillation, n (%)	25 (33.8)	11 (42.3)	0.44
Number of prescriptions, median (IQR)	8.0 (5.3-11.0)	8.0 (6.0-10.5)	0.78
Logistic EuroSCORE I, median (IQR)	15.0 (10.0-19.0)	16.5 (11.0-21.9)	0.42
Left ventricular ejection fraction in %, median (IQR)	55.0 (45.0-55.0)	52.5 (44.3-55.0)	0.73
eGFR, mean ml/min/1.73m ² (SD)	55.9 (15.8)	55.7 (18.9)	0.94

SD= Standard Deviation, ADL= Activities of Daily Living, iADL= Instrumental Activities of Daily Living, IQR= Interquartile Range, MMSE= Mini Mental State Examination