

Supplementary materials




Figure S1. Receiver Operating Characteristic (ROC) curve for International Council for Standardization in Haematology (ICSH) method of schistocyte quantitation for venom-induced consumption coagulopathy (VICC) with acute kidney injury (AKI) versus VICC without AKI

Table S1. Sensitivity and specificity for International Council for Standardization in Haematology method of schistocyte quantitation for venom induced consumption coagulopathy (VICC) with acute kidney injury (AKI) versus VICC without AKI

Schistocytes %	Sensitivity %	95% CI	Specificity %	95% CI	Likelihood ratio
> 0.05	100	83.2% - 100%	3.85	1.51% - 9.47%	1.04
> 0.15	100	83.2% - 100%	13.5	8.19% - 21.3%	1.16
> 0.25	100	83.2% - 100%	23.1	16.0% - 32.0%	1.30
> 0.35	100	83.2% - 100%	31.7	23.6% - 41.2%	1.46
> 0.45	89.5	68.6% - 98.1%	38.5	29.7% - 48.1%	1.45
> 0.55	89.5	68.6% - 98.1%	48.1	38.7% - 57.6%	1.72
> 0.65	89.5	68.6% - 98.1%	51.0	41.5% - 60.4%	1.82
> 0.75	89.5	68.6% - 98.1%	55.8	46.2% - 64.9%	2.02
> 0.85	89.5	68.6% - 98.1%	59.6	50.0% - 68.5%	2.22
> 0.95	89.5	68.6% - 98.1%	71.2	61.8% - 79.0%	3.10
> 1.05	89.5	68.6% - 98.1%	76.0	66.9% - 83.2%	3.72
> 1.15	89.5	68.6% - 98.1%	78.8	70.0% - 85.6%	4.23
> 1.25	84.2	62.4% - 94.5%	82.7	74.3% - 88.8%	4.87
> 1.35	78.9	56.7% - 91.5%	84.6	76.5% - 90.3%	5.13
> 1.45	78.9	56.7% - 91.5%	86.5	78.7% - 91.8%	5.86
> 1.55	73.7	51.2% - 88.2%	92.3	85.6% - 96.1%	9.58
> 1.65	68.4	46.0% - 84.6%	94.2	88.0% - 97.3%	11.9
> 1.75	63.2	41.0% - 80.9%	97.1	91.9% - 99.2%	21.9
> 2.00	57.9	36.3% - 76.9%	99.0	94.8% - 100%	60.2
> 2.35	52.6	31.7% - 72.7%	99.0	94.8% - 100%	54.7
> 2.75	42.1	23.1% - 63.7%	99.0	94.8% - 100%	43.8
> 3.10	36.8	19.1% - 59.0%	99.0	94.8% - 100%	38.3
> 3.65	31.6	15.4% - 54.0%	100	96.4% - 100%	
> 4.80	26.3	11.8% - 48.8%	100	96.4% - 100%	
> 5.55	21.1	8.51% - 43.3%	100	96.4% - 100%	
> 6.75	15.8	5.52% - 37.6%	100	96.4% - 100%	
> 9.50	10.5	1.87% - 31.4%	100	96.4% - 100%	
> 11.3	5.26	0.270% - 24.6%	100	96.4% - 100%	