

Additional File 5. Comparison of baseline study characteristics between complete (n=6,691 and n=4,624) and incomplete (n=11,920) cases

| % or Mean (SD) for measures at baseline (2002/2005) | Complete cases* | Incomplete cases | P-value of sample differences** |
|---|-----------------|------------------|---------------------------------|
| Immediate recall (0-30) | 20.8 (4.3) | 20.3 (4.4) | P<0.001 |
| Delayed recall (0-10) | 7.0 (1.9) | 6.8 (2.0) | P<0.001 |
| Verbal fluency | 20.6 (6.9) | 20.0 (6.8) | P<0.001 |
| Processing speed (0-65) | 17.7 (5.5) | 17.3 (5.5) | P<0.001 |
| Country | | | P<0.001 |
| Czech Republic | 30.9% | 30.2% | |
| Poland | 34.0% | 33.8% | |
| Russia | 35.1% | 36.0% | |
| Age (years) | 62.2 (6.0) | 62.4 (5.9) | P<0.001 |
| Female | 53.7% | 53.0% | P<0.001 |
| Network size of friends | | | P<0.001 |
| None | 46.5% | 50.6% | |
| 1 or 2 | 35.3% | 32.8% | |
| 3 to 5 | 14.3% | 13.0% | |
| More than 5 | 4.0% | 3.5% | |
| Network size of relatives | | | |
| None | 40.5% | 43.6% | |
| 1 or 2 | 35.3% | 34.7% | |
| 3 to 5 | 20.6% | 18.5% | |
| More than 5 | 3.6% | 3.3% | |
| Contact frequency with friends | | | P<0.001 |
| No friends | 6.7% | 8.0% | |
| Less than once a month | 26.7% | 29.1% | |
| About once a month | 21.7% | 20.4% | |
| Several times a month | 16.3% | 15.2% | |
| About once a week | 16.4% | 15.6% | |
| Several times a week | 12.2% | 11.8% | |
| Contact frequency with relatives | | | |
| No relatives | 2.6% | 2.9% | |
| Less than once a month | 18.6% | 21.1% | |
| About once a month | 14.1% | 14.3% | |
| Several times a month | 14.5% | 13.8% | |
| About once a week | 24.5% | 23.4% | |
| Several times a week | 25.8% | 24.6% | |
| Participation in social activities | | | P<0.001 |
| Never or not a member | 81.8% | 84.3% | |
| At least several times a year | 8.8% | 7.7% | |
| Several times a month or more | 9.4% | 8.0% | |
| Educational level | | | P<0.001 |
| Primary or lower | 9.6% | 13.4% | |
| Vocational | 26.2% | 26.7% | |
| Secondary | 37.0% | 35.8% | |
| University | 27.3% | 24.1% | |
| Number of household amenities (0-12) | 6.3 (2.2) | 6.1(2.2) | P<0.001 |
| Not working | 64.6% | 67.1% | P<0.001 |
| Not partnered | 25.6% | 27.1% | P<0.001 |
| Smoking status | | | P<0.001 |
| Current | 20.4% | 22.7% | |
| Former | 25.2% | 25.2% | |
| Never | 54.3% | 52.1% | |
| Alcohol drinking frequency | | | P<0.001 |
| Never | 21.5% | 24.0% | |
| <1/month | 29.9% | 30.0% | |
| 1-3/month | 20.3% | 19.4% | |
| 1-4/week | 21.9% | 20.6% | |
| More than 5/week | 6.4% | 6.0% | |
| Alcohol intake per week (grams) | 2580.7 (6134.5) | 2544.4 (6386.9) | P<0.001 |

| % or Mean (SD) for measures at baseline (2002/2005) | Complete cases* | Incomplete cases | P-value of sample differences** |
|--|------------------------|-------------------------|--|
| Physical activity per week (hours) | 21.2 (14.3) | 20.1 (14.2) | P<0.001 |
| Self-rated health | | | P<0.001 |
| Very poor or poor | 14.5% | 16.9% | |
| Fair | 59.6% | 59.4% | |
| Very good or good | 26.0% | 23.7% | |
| Number of chronic diseases (0-4) | 3.7 (0.6) | 3.6 (0.7) | P<0.001 |
| CES-D 20 score (0-60) | 10.9 (7.9) | 11.1 (8.1) | P<0.001 |

Note: Covariate baseline data on age, sex, and country were complete for all participants.

*Complete case estimates are based on the analytic sample for frequency of contact with social network members and participation in social activities (n=6,691), with exception to baseline frequencies on social network size which was calculated using the sub-sample of participants with these data (n=4,624).

**Proportion and mean differences between complete and incomplete cases were compared using chi-squared tests and t-tests.