

Supplemental Table S1. Sensitivity Analysis Excluding Subjects with the Occurrence of End Points within 1 Year of Follow-up (Center Column) or Excluding Subjects with a History of Cancer at Baseline (Right Column)

IFG exposure score	Excluding subjects with the occurrence of end points within 1 year (1-year lag period)		Excluding subjects with a history of cancer at baseline	
	Model 1	Model 2	Model 1	Model 2
All-cause mortality				
0	1 (reference)	1 (reference)	1 (reference)	1 (reference)
1	1.11 (1.07–1.14)	1.10 (1.06–1.13)	1.10 (1.07–1.14)	1.09 (1.06–1.13)
2	1.15 (1.10–1.19)	1.13 (1.09–1.17)	1.15 (1.11–1.20)	1.13 (1.09–1.17)
3	1.21 (1.15–1.27)	1.18 (1.12–1.24)	1.20 (1.15–1.26)	1.17 (1.12–1.23)
4	1.18 (1.11–1.26)	1.15 (1.08–1.23)	1.18 (1.11–1.26)	1.15 (1.07–1.22)
Myocardial infarction				
0	1 (reference)	1 (reference)	1 (reference)	1 (reference)
1	1.09 (1.05–1.14)	1.07 (1.02–1.12)	1.09 (1.05–1.14)	1.07 (1.02–1.12)
2	1.09 (1.04–1.15)	1.05 (1.00–1.11)	1.10 (1.05–1.16)	1.06 (1.01–1.11)
3	1.08 (1.01–1.15)	1.02 (0.96–1.09)	1.10 (1.03–1.17)	1.04 (0.97–1.10)
4	1.05 (0.95–1.15)	0.98 (0.89–1.08)	1.05 (0.96–1.15)	0.98 (0.89–1.07)
Stroke				
0	1 (reference)	1 (reference)	1 (reference)	1 (reference)
1	1.09 (1.04–1.14)	1.07 (1.02–1.12)	1.10 (1.05–1.15)	1.08 (1.03–1.12)
2	1.11 (1.05–1.17)	1.07 (1.01–1.13)	1.13 (1.07–1.19)	1.08 (1.03–1.14)
3	1.12 (1.04–1.20)	1.06 (0.99–1.14)	1.11 (1.04–1.19)	1.05 (0.98–1.12)
4	1.11 (1.01–1.22)	1.05 (0.95–1.15)	1.13 (1.04–1.24)	1.06 (0.97–1.16)

Hazard ratios and 95% confidence intervals of all-cause mortality, myocardial infarction (MI), and stroke according to impaired fasting glucose IFG exposure score. Model 1: Adjusted for age, sex, alcohol consumption, smoking, physical activity, and body mass index; Model 2: Adjusted for age, sex, alcohol consumption, smoking, physical activity, body mass index, hypertension, and dyslipidemia.