
SARS-CoV-2 IgG seropositivity in a cohort of 449 non-hospitalized individuals during Spanish COVID-19 lockdown.

Patricia Torres Martínez MD, Paula Diaque García, María Rubio Salas, Raquel Rodríguez Sánchez MD, José Avendaño-Ortíz PhD, Sandra Guerrero-Monjo PhD, Felipe García, Miguel Ángel Llamas PhD, Eduardo López-Collazo PhD, Paula Saz-Leal PhD, Carlos del Fresno PhD



Supplementary Figure 1. Symptomatology-dependent seropositivity of SARS-CoV-2 within age ranges. Dots represent IgG+ symptomatic (blue) and asymptomatic (grey) subjects at given age ranges. Dots and vertical lines represent frequency and 95% confidence interval, respectively. Frequencies are further detailed in the bottom table.

Supplementary Table 1. COVID-19 compatible symptoms and close contact of study participants by occupation.

		Frequency			Univariate analysis	
		%	(95% CI)	OR	95% CI	P-value
Symptoms compatible with COVID-19 ¹	Others ²	70.47%	(65.46-75.15)	1		0.366 ³
	Healthcare workers	72.09%	(56.33-84.67)	1.08	0.53 to 2.19	
	Firefighters	66.67%	(40.99-86.66)	0.84	0.31 to 2.29	
	Police/Public safety	55.17%	(35.69-73.55)	0.52	0.24 to 1.11	
Any contact with confirmed case	Others ²	57.10%	(51.80-62.28)	1		- - <0.001 ³
	Healthcare workers	79.07%	(63.96-89.96)	2.83	1.32 to 6.09	
	Firefighters	94.44%	(72.71-99.86)	12.77	1.68 to 97.05	
	Police/Public safety	72.41%	(52.76-87.27)	1.97	0.85 to 4.57	

¹ Fever, headache, cough, odynophagia, asthenia, myalgia, ageusia, anosmia, dyspnea, , gastrointestinal symptoms cutaneous manifestations and/or pneumonia diagnosis.

OR: Odds Ratio. 95% CI: 95% Confidence Interval.

Supplementary Table 2. Forward logistic regression model.

		Forward multivariate analysis				
		OR	95% CI	P-value		
Smoker	No	1				
Smoker	Yes	0.44	0.20 to 0.95	0.037		
	Never done	1				
PCR estatus	Negative	1.42	0.58 to 3.48	0.442		
	Positive	25.12	2.62 to 240.95	0.005		
Contact with	No contact	1				
confirmed case	Household member	3.72	2.03 to 6.79	<0.001		
commined case	Non-cohabitating	1.48	0.79 to 2.78	0.216		
COVID-19	No	1				
Symptoms	Yes	2.49	1.33 to 4.66	0.004		
Type of	Ageusia/Anosmia	13.67	7.32 to 25.53	<0.001		
symptoms	Odynophagia	0.36	0.19 to 0.66	0.001		

OR: Odds Ratio. 95% CI: 95% Confidence Interval.

Variables significantly associated with seropositivity are highlighted in bold.

² Those patients that are not among the other three specific professions.

³ Chi-square test.