

Supplemental Table 1. Selected baseline comorbidities and migraine characteristics by provider specialty

	All patients N = 9753	Provider specialty	
		Neurologist N = 7067	Primary care <sup>a</sup> N = 1387
Comorbidities, n (%)			
Asthma	727 (7.5)	560 (7.9)	75 (5.4)
Hypertension	1790 (18.4)	1323 (18.7)	215 (15.5)
Acute myocardial infarction	37 (0.4)	24 (0.3)	5 (0.4)
Ischemic stroke	77 (0.8)	69 (1.0)	3 (0.2)
Constipation	306 (3.1)	244 (3.5)	17 (1.2)
Irritable bowel syndrome	232 (2.4)	185 (2.6)	21 (1.5)
Complications of constipation	187 (1.9)	141 (2.0)	8 (0.6)
Anxiety	2125 (21.8)	1589 (22.5)	260 (18.7)
Depression	1606 (16.5)	1212 (17.2)	193 (13.9)
Other (non-migraine) headaches	2829 (29.0)	2277 (32.2)	264 (19.0)
Low back pain	1511 (15.5)	1144 (16.2)	132 (9.5)
Osteoarthritis/spondylosis	1411 (14.5)	1085 (15.4)	105 (7.6)
Fibromyalgia	549 (5.6)	430 (6.1)	48 (3.5)
No comorbidities			
Migraine characteristics			
Migraine with aura, including persistent aura	2611 (26.8)	1937 (27.4)	361 (26.0)
Migraine with aura, excluding persistent aura	2530 (25.9)	1878 (26.6)	347 (25.0)
Chronic migraine	6274 (64.3)	5009 (70.9)	559 (40.3)
Menstrual migraine	259 (2.7)	195 (2.8)	35 (2.5)
Status migrainosus	2607 (26.7)	2053 (29.1)	288 (20.8)

<sup>a</sup>Family practice/internal medicine physician and no neurologist or pain specialist/anesthesiologist