

Supplemental information

ELAVL4, splicing, and glutamatergic dysfunction precede neuron loss in *MAPT* mutation cerebral organoids

Kathryn R. Bowles, M. Catarina Silva, Kristen Whitney, Taylor Bertucci, Joshua E. Berlind, Jesse D. Lai, Jacob C. Garza, Nathan C. Boles, Sidhartha Mahali, Kevin H. Strang, Jacob A. Marsh, Cynthia Chen, Derian A. Pugh, Yiyuan Liu, Ronald E. Gordon, Susan K. Goderie, Rebecca Chowdhury, Steven Lotz, Keith Lane, John F. Crary, Stephen J. Haggarty, Celeste M. Karch, Justin K. Ichida, Alison M. Goate, and Sally Temple

Table S1. Human iPSC lines used to generate 3D cerebral organoids.

<i>MAPT</i>	Age at Biopsy	iPSC Line	Abbreviated ID	Sanger seq. confirmed?	Karyotype
<i>V337M</i>	40s	ND32951A.15Δ2B09	NDB09	yes	normal
<i>V337V</i>	40s	ND32951A.15Δ1B06	NDB06	yes	normal
<i>V337M</i>	60s	GIH-6-C1-ΔA02	GIH6-A02	yes	normal
<i>V337V</i>	60s	GIH-6-C1-Δ1E11	GIH6-E11	yes	normal
<i>V337M</i>	30s	GIH-7-C2-Δ2A01	GIH7-A01	yes	normal
<i>V337V</i>	30s	GIH-7-C2-Δ2B12	GIH7-B12	yes	normal
<i>V337V</i>	30s	GIH-7-C2-Δ2F02	GIH7-F02	yes	normal
<i>WT</i>	50s	ND03231	3231	no	normal