

Supplementary table 1. Distribution of deaths among COVID-19 positive patients in selected hospitals of Bangladesh, June-August 2020.

	Hospital death n= 16	Home death n=5
Age (years)		
Lowest, highest	40, 85	51, 90
Median (IQR)	65 (55-69)	64 (52-70)
Sex		
Male	11 (69)	3 (60)
Timing of death from symptom onset		
0-7 days	6 (38)	6 (38) Cum.
8-15 days	9 (56)	15 (94)
16-40 days	1 (6)	16 (100)
Manner of death		
Disease	16	5 (100)
Accident	0	0
Sudden death (heart attack)	0	0
Comorbidity present (anyone)		
DM	15 (94%)	4 (80)
Asthma/COPD	9 (56)	2 (40)
Asthma/COPD	4 (25)	2 (40)
Heart disease/HTN	6 (37)	2 (40)
Chronic renal disease	4 (25)	0
Cancer (uterine)	4 (25)	0
Clinical course of treatment		
Oxygen required	16 (100)	3 (60) [2 got oxygen at home]
ICU admitted	3 (19)	
CCU admitted	1 (6)	
Dialysis required	1 (6)	
Cause of death as stated from hospital as reported		
COVID-19 + Respiratory failure	14 (88)	
Pneumonia	1 (6)	
Chronic renal failure	1 (6)	