## Supplementary Figure 1: Tools to measure severity of bleeding

## Rectal bleeding score

Symptoms	Score	Diarrhoea			
Diarrhoea	0	Absent			
	1	1-3 stools/24 hours			
	2	4-6 stools/24 hours			
	3	>6 stools/24 hours			
Bleeding	0 No blood				
	1	Blood on toilet paper/in stool			
	2	Blood in toilet bowl			
	3	Heavy bleeding with clots			
	4	Bleeding requiring transfusion			
Tenesmus/rectal	0	Absent			
pain	1	Mild tenesmus not requiring any drug			
	2	Tenesmus requiring analgesics/antispasmodics			
	3	Severe tenesmus requiring daily use of analgesics/antispasmodics			

## Endoscopy severity score

Grade	Score	Description
Normal	0	Normal mucosa
Mild	3	Erythema and/or telangectasia
Moderate	6	Friability
Severe	9	Ulceration and/or necrosis

Total score: \_\_\_\_\_

Patient-reported diary

Day	Time	Stool passed Yes/No	Stool consistency (stool chart)	Blood on wiping Yes/No	Blood in pan Yes/No	Is the bleeding less heavy Yes/No	Clots in pan Yes/No	Accidents with blood Yes/No
1								
2								
_								

## Supplementary Table 2: Zinicola Score

	Distribution of telangectasia	Surface area of telangectasia	Presence of fresh blood
0			None
1	Distal rectum (within 10cm anal verge)	<50%	Some
2	Entire rectum	>50%	

2-Grade A; 3-Grade B; 4/5-Grade C

Grade \_\_\_\_\_