

Supplementary

Table S1 Endpoints in PP

Variable	Value
Patients with complete follow-up (n, %)	
With indication for circumcision	34 (47.9%)
Without indication for circumcision	37 (52.1%)
“Are you satisfied by the results?” (n, %)	
Yes	41 (57.7%)
No	25 (35.2%)
I don’t know	5 (7.0%)
“Would you recommend it?” (n, %)	
Yes	57 (80.3%)
No	9 (12.7%)
I don’t know	5 (7.0%)
Penile sensitivity (n, %)	
Reduced	13 (18.3%)
Increased	12 (16.9%)
Not modified	46 (64.8%)
Erectile function (n, %)	
Improved	10 (14.1%)
Worsened	5 (7.0%)
Not modified	56 (78.9%)
Ejaculation time (n, %)	
Improved	9 (12.7%)
Worsened	1 (1.4%)
Not modified	61 (85.9%)
Orgasmic function (n, %)	
Improved	1 (1.4%)
Worsened	7 (9.9%)
Not modified	63 (88.7%)
Persistent painful sexual intercourse (n, %)	
Yes	17 (23.9%)
No	28 (39.4%)
I don’t know	26 (36.6%)
Good cosmetic and functional result (n, %)	44 (62.0%)
Persistent phimosis (n, %)	41 (57.7%)