



Supplemental Figures


1) Forest Plot of Pooled prevalence of Fatigue in Post COVID-19 patients.




2) Forest Plot of Pooled prevalence of Dyspnea in Post COVID-19 patients.




3) Forest Plot of Pooled prevalence of Cough in Post COVID-19 patients.




4) Forest Plot of Pooled prevalence of Anosmia in Post COVID-19 patients.




5) Forest Plot of Pooled prevalence of Sleep Disturbances in Post COVID-19 patients.




6) Forest Plot of Pooled prevalence of Chest Pain in Post COVID-19 patients.




7) Forest Plot of Pooled prevalence of Headache in Post COVID-19 patients.




8) Forest Plot of Pooled prevalence of Arthralgia in Post COVID-19 patients.




9) Forest Plot of Pooled prevalence of Mental health/PTSD in Post COVID-19 patients.



10) Meta-regression between poor quality of life (log-event) and anosmia in Post COVID-19 patients.



11) Meta-regression between poor quality of life (log-event) and Dyspnea in Post COVID-19 patients.



Supplemental: Risk of bias of included studies

Author, Country	Newcastle–Ottawa Scale			Overall risk of bias
	Selection	Comparability	Exposure	
Chopra et al., USA	**	*	*	High
Carfi et al., Italy	***	*	**	Moderate
Haplin et al., UK	**	*	*	High
Taboada et al., Spain	**	*	*	High
Jacobs et al., USA	***	*	**	Low
Tabacof et al., USA	***	*	**	Low
Valent et al., France	****	*	*	Moderate
Arab-Zozani et al., Iran	****	*	**	Moderate
Mandal et al., UK	***	*	**	Low
Huang et al., China	***	*	**	Moderate
Garrigues et al., France	**	*	*	High
Moreno-Pérez et al., Spain	****	*	**	Moderate