

Supplemental Table 3. Intraoperative antibiotics for intraabdominal surgery patients, by 30-day mortality status.

Antibiotic	Total 4322	Died 62 (1.4)	Alive 4260 (99)	P-Value
Ampicillin	20 (0.5)	0 (0.0)	20 (0.5)	1.0
Aztreonam	6 (0.1)	0 (0.0)	6 (0.1)	1.0
Bactrim	1 (0.0)	0 (0.0)	1 (0.0)	1.0
Cefamandole	2 (0.0)	0 (0.0)	2 (0.0)	1.0
Cefazolin	1701 (39)	9 (15)	1692 (40)	<0.0001
Cefepime	6 (0.1)	1 (1.6)	5 (0.1)	0.15
Cefotaxime	1 (0.0)	0 (0.0)	1 (0.0)	1.0
Cefoxitin	272 (6.3)	7 (11)	265 (6.2)	0.2
Ceftazidime	1 (0.0)	0 (0.0)	1 (0.0)	1.0
Ceftriaxone	3 (0.1)	0 (0.0)	3 (0.1)	1.0
Clindamycin	167 (3.9)	4 (6.5)	163 (3.8)	0.5
Fluconazole	22 (0.5)	1 (1.6)	21 (0.5)	0.7
Gentamicin	26 (0.6)	1 (1.6)	25 (0.6)	0.8
Levofloxacin	242 (5.6)	2 (3.2)	240 (5.6)	0.6
Linezolid	1 (0.0)	0 (0.0)	1 (0.0)	1.0
Meropenem	5 (0.1)	1 (1.6)	4 (0.1)	0.11
Metronidazole	444 (10)	13 (21)	431 (10)	0.01
Piperacillin	14 (0.3)	0 (0.0)	14 (0.3)	1.0
Tobramycin	1 (0.0)	0 (0.0)	1 (0.0)	1.0
Unasyn	575 (13)	5 (8.1)	570 (13)	0.3
Vancomycin	52 (1.2)	1 (1.6)	51 (1.2)	1.0
Zosyn	519 (12)	10 (16)	509 (12)	0.4

Counts and (%) are displayed for each group and compared with the chi-square test.