Supplementary Material Table 2. Antimicrobial Prevalence by Surgical Prophylaxis Site for Adult, Pediatric and Neonatal Wards

Adult	All	AMW	НО-	T-	P-	ASW	AICU
	wards		AMW	AMW	AMW		
Total number of antimicrobials prescribed for surgical prophylaxis, N	567	86	4	0	1	403	73
Prophylaxis site, N							
General surgical prophylaxis without targeting a specific site	11	0	4	0	0	7	0
Other	2	0	0	0	0	0	2
Bone and Joint	176	10	0	0	0	158	8
Central Nervous System	29	4	0	0	0	14	11
Cardiovascular System	107	19	0	0	0	44	44
Ear, Nose, Throat	31	3	0	0	0	26	2
Gastrointestinal Tract	78	6	0	0	0	67	5
Obstetrics or Gynecological	68	38	0	0	0	30	0
Respiratory Tract	7	0	0	0	1	6	0
Urinary Tract	58	6	0	0	0	51	1
Pediatric and Neonatal	All wards	PMW	GNMW	HO- PMW	PSW	PICU	NICU
Total number of antimicrobials prescribed for surgical prophylaxis,	11	3	0	0	5	0	3
Prophylaxis site, N							
Bone and Joint	2	0	0	0	2	0	0
Ear, Nose, Throat	1	0	0	0	1	0	0
Gastrointestinal Tract	8	3	0	0	2	0	3

Abbreviations: AMW, adult medical ward; HO-AMW, hematology-oncology-AMW, T-AMW, transplant-AMW; P-AMW, pneumology-AMW; ASW, adult surgical ward; AICU, adult intensive care unit; PMW, pediatric medical ward; GNMW, general neonatal medical ward; PSW, pediatric surgical ward; PICU, pediatric intensive care unit; NICU, neonatal intensive care unit