

Patient Characteristic		Quartile of Neighborhood Disadvantage Index				p-value
		Quartile 1 (Least Disadvantaged)	Quartile 2	Quartile 3	Quartile 4 (Most Disadvantaged)	
	Overall	<0.73	0.73 to <1.28	1.28 to <1.96	≥1.96	
Age at diagnosis, median (IQR)	29.3 (6.9)	32.0 (27.0-36.0)	31.0 (27.0-34.0)	27 (21.8-31.5)	28.0 (23.0-32.0)	0.002
Black/African American race, n (%)	121 (55.0)	10 (16.1)	22 (41.6)	42 (80.8)	47 (88.7)	<0.001
Pre-pregnancy hypertension, n (%)	80 (36.4)	23 (37.1)	14 (26.4)	22 (42.3)	21 (39.6)	0.345
Twin pregnancy, n(%)	28 (12.7)	8 (12.9)	12 (22.6)	3 (5.8)	5 (9.4)	0.058
Day of diagnosis, median (IQR)	7.0 (1.0-28.0)	6.0 (2.0-15.0)	5.0 (1.0-26.0)	7.0 (2.0-38.0)	15.5 (2.0-35.0)	0.397
Timing of diagnosis						0.365
Prepartum	24 (10.9)	5 (8.1)	6 (11.3)	6 (11.5)	7 (13.2)	
<30 days postpartum	150 (68.2)	49 (79.0)	37 (69.8)	33 (63.5)	31 (58.5)	
≥30 days postpartum	46 (20.9)	8 (12.9)	10 (18.9)	13 (25.0)	15 (28.3)	
Lowest ejection fraction, median (IQR)	22.5 (15.0-35.0)	30 (17.5-37.5)	26.3 (15.0-35.0)	20.0 (12.5-30.0)	17.5 (12.5-37.5)	0.270
Patient Characteristic		Quartile of Median Household Income for Block Group				p-value
		Quartile 4 (Least Disadvantaged)	Quartile 3	Quartile 2	Quartile 1 (Most Disadvantaged)	
		Overall	≥\$75,649	\$46,266 to <\$75,649	\$26,466 to <\$46,266	
Age at diagnosis, median (IQR)	29.3 (6.9)	33.0 (30.0-36.0)	30.0 (25.0-34.0)	29.0 (22.0-32.0)	27.5 (23.0-31.0)	<0.001
Black/African American race, n (%)	121 (55.0)	6 (11.1)	22 (40.0)	43 (75.4)	50 (92.6)	<0.001
Pre-pregnancy hypertension, n (%)	80 (36.4)	22 (40.7)	16 (29.1)	26 (45.6)	16 (29.6)	0.182
Twin pregnancy, n(%)	28 (12.7)	10 (18.5)	10 (18.2)	4 (7.0)	4 (7.4)	0.104
Day of diagnosis, median (IQR)	7.0 (1.0-28.0)	5.0 (1.0-12.0)	9.0 (2.0-61.0)	7.0 (1.0-24.0)	7 (0.5-36.5)	0.320
Timing of diagnosis						0.125
Prepartum	24 (10.9)	4 (7.4)	3 (5.5)	9 (15.8)	8 (14.8)	
<30 days postpartum	150 (68.2)	42 (77.8)	39 (70.9)	39 (68.4)	30 (55.6)	
≥30 days postpartum	46 (20.9)	8 (14.8)	13 (223.6)	9 (15.8)	16 (29.6)	
Lowest ejection fraction, median (IQR)	22.5 (15.0-35.0)	35.0 (25.0-40.0)	20.2 (12.5-33.8)	20.0 (12.5-35.0)	17.5 (12.5-32.5)	0.005
Patient Characteristic		Quartile of Proportion of Residents with Lower than High School Education for Block Group				p-value
		Quartile 1	Quartile 2	Quartile 3	Quartile 4	
		Overall	<5.4%	5.4% to <12.1%	12.1% to <22.3%	
Age at diagnosis, median (IQR)	29.3 (6.9)	31.0 (28.0-35.0)	31.0 (26.5-33.5)	28.0 (24.0-32.0)	27.0 (22.0-34.0)	0.006
Black/African American race, n (%)	121 (55.0)	13 (23.6)	19 (33.9)	41 (74.6)	48 (88.9)	<0.001
Pre-pregnancy hypertension, n (%)	80 (36.4)	19 (35.6)	19 (33.9)	21 (38.2)	21 (38.9)	0.929
Twin pregnancy, n(%)	28 (12.7)	7 (12.7)	14 (25.0)	4 (7.3)	3 (5.6)	0.009
Day of diagnosis, median (IQR)	7.0 (1.0-28.0)	5.0 (1.0-12.0)	7.0 (1.0-26.0)	8.0 (2.0-45.0)	9.0 (1.0-35.0)	0.506
Timing of diagnosis						0.614
Prepartum	24 (10.9)	7 (12.7)	4 (7.1)	6 (10.9)	7 (13.0)	
<30 days postpartum	150 (68.2)	40 (72.7)	41 (73.2)	37 (67.3)	32 (59.3)	
≥30 days postpartum	46 (20.9)	8 (14.6)	11 (19.6)	12 (21.8)	15 (27.8)	
Lowest ejection fraction, median (IQR)	22.5 (15.0-35.0)	31.3 (20.0-37.5)	27.5 (12.5-37.0)	20.0 (10.0-33.8)	17.5 (12.5-35.0)	0.123