

	Survived (n = 48)	Died (n = 24)	P-value
Median Age (IQR)	57.0 (50.0 – 62.5)	60.0 (59.0 – 65.0)	0.064
Male (%)	36 (75.0)	18 (75.0)	
NIDDM (%)	9 (18.8)	10 (41.7)	0.038
IDDM (%)	4 (8.3)	3 (12.5)	0.574
Mean BMI (SD)	29.5 (5.7)	28.2 (4.8)	0.321
Admission HbA1c (SD)	50.2 (17.0)	60.8 (34.7)	0.284
Admission Vitamin D (SD)	31.7 (16.8)	22.7 (10.1)	0.116
Insulin Therapy (%)	21 (43.8)	15 (62.5)	0.134
Steroid Therapy (%)	8 (16.7)	11 (45.8)	0.008
Antibiotic Therapy (%)	37 (77.1)	22 (91.7)	0.129
Pabrinex Therapy (%)	14 (29.2)	8 (33.3)	0.160
Vitamin D Supplementation (%)	3 (6.3)	4 (16.7)	0.164
Median ITU LOS (IQR)	25.0 (9.0 – 38.8)	13.5 (6.0 – 31.5)	0.175
Median ventilator days (IQR)	20.0 (12.5 – 34.3)	12.0 (6.0 – 30.3)	0.190
Median RRT days (IQR)	0.0 (0.0 – 9.0)	5.0 (0.0 – 10.5)	0.276