

**Supplementary Table 13. Spearman correlation analysis of anti-spike IgG levels at t4 versus IgG levels at earlier collection time points** (data for t4 vs t1 presented in Figure 7B).

Correlation analysis (n=116)			
IgG OD norm	Age Group	p-value*	$\rho^\dagger$
t4 vs t1	All	$5.623 \times 10^{-12}$	0.5845
	27-70	$7.993 \times 10^{-07}$	0.6213
	71-93	$2.193 \times 10^{-05}$	0.5214
t4 vs t2	All	$5.749 \times 10^{-09}$	0.5082
	27-70	$3.593 \times 10^{-05}$	0.5334
	71-93	$8.605 \times 10^{-05}$	0.4866
t4 vs t3	All	$<2.2 \times 10^{-16}$	0.9426
	27-70	$<2.2 \times 10^{-16}$	0.9331
	71-93	$<2.2 \times 10^{-16}$	0.9444

t0 – sera collected on day of 1<sup>st</sup> vaccine dose; t1 – sera collected 21 days post-1<sup>st</sup> vaccine dose; t2 – sera collected 42 days post-1<sup>st</sup> vaccine dose; t3 - sera collected ~140 days post-1<sup>st</sup> vaccine dose; t4 - sera collected 180 days post-1<sup>st</sup> vaccine dose.

\* Spearman correlation p-value

† Spearman correlation coefficient