

1 **Table E1. Adjusted associations Between Participant Characteristics and Timing of**
 2 **Introduction of specific allergen:**

	Peanut	Milk	Egg
Participant Characteristics Associated with: Early Introduction of specific allergen (6 months or before)	Odds Ratio (95% CI)	Odds Ratio (95% CI)	Odds Ratio (95% CI)
White (vs Black)	2.6 (1.1-7.2)*	2.7 (1.1-6.7)*	3.3 (1.0-14.9)
Age (continuous--in years)	0.8 (0.7-0.9)*	0.9 (0.8-1.0)	0.8 (0.7-1.0)*
Multiple (vs single) Food Allergies	1.5 (0.6-3.7)	0.8 (0.3-2.5)	0.8 (0.7-1.0)
Female (vs Male)	1.9 (0.9-3.9)	1.0 (0.4-2.5)	1.3 (0.5-3.1)
Eczema (vs no eczema)	1.0 (0.4-2.8)	4.7 (1.4-19.5)*	0.6 (0.2-1.7)
Patient Characteristics Associated with: Later introduction of specific allergen (after 11 months of age)			
White (vs Black)	0.3 (0.1-0.5)*	0.2 (0.1-0.6)*	0.5 (0.2-1.2)
Age (continuous--in years)	1.3 (1.2-1.5)*	1.1 (1.0-1.2)	1.3 (1.1-1.5)
Multiple (vs single) Food Allergies	0.9 (0.5-1.8)	1.6 (0.5-6.0)	3.0 (1.0-10.8)
Female (vs Male)	0.6 (0.4-1.2)	1.1 (0.4-2.6)	0.6 (0.3-1.4)
Eczema (vs no eczema)	0.8 (0.4-1.9)	0.4 (0.1-1.2)	2.1 (0.7-6.5)

3 CI: Confidence Interval

4 Note: Model also includes fixed effects adjustment for study recruitment site

5 * Variable is a statistically significant predictor of model-specific food allergen introduction
 6 timing at a two-sided p<.05

7

Figure E1: Racial Differences in Reported Timing of Peanut Introduction by child race and birth-year [Estimated % (95% Confidence Interval)]





Figure E1 Legend: Racial Differences in Reported Timing of Peanut Introduction by child race and birth-year [Estimated % (95% Confidence Interval)]

Journal Pre-proof