

Supplemental Table 2. Cases of Postoperative Endophthalmitis Within 2 weeks following Cataract Surgery Diagnosed with or without Supporting Clinical Findings and Corresponding Odds Ratios of Developing Endophthalmitis by Surgery Group.

		Unilateral N=1,712,590 ¹	DSBCS N=3,695,440 ¹	DSBCS and Unilateral N=5,408,030 ¹	ISBCS N=165,609 ¹	Total N=5,573,639 ¹
Endophthalmitis Diagnosis	Either Eye ²	1,094 (0.064%)	1,915 (0.052%)	3,009 (0.056%)	74 (0.045%)	3,083 (0.055%)
	First Eye	1,094 (0.064%)	538 (0.015%)	1,632 (0.030%)	39 (0.024%)	1,671 (0.030%)
	Second Eye ³	-	1,393 (0.038%)	1,393 (0.038%)	38 (0.023%)	1,431 (0.037%)
	Both Eyes ³	-	16 (0.0004%)	16 (0.0004%)	3 (0.002%)	19 (0.0005%)
Diagnosis with Supporting Clinical Findings	Either Eye ²	941 (0.055%)	1,595 (0.043%)	2,536 (0.047%)	61 (0.037%)	2,597 (0.047%)
	First Eye	941 (0.055%)	425 (0.012%)	1,366 (0.025%)	31 (0.019%)	1,397 (0.025%)
	Second Eye ³	-	1,176 (0.032%)	1,176 (0.032%)	30 (0.018%)	1,206 (0.031%)
	Both Eyes ³	-	6 (0.0002%)	6 (0.0002%)	0 (0.0%)	6 (0.0002%)

¹Statistics presented: n (%)

²Patients had postoperative endophthalmitis in either their first and/or second surgery eye.

³Percentages taken out of the number of participants who had bilateral cataract surgery (n = 3,695,440, 165,609, and 3,861,049)

DSBCS, delayed sequential bilateral cataract surgery; ISBCS, immediate sequential bilateral cataract surgery; CI, confidence intervals