

Higher Ambient Nitrogen Dioxide Is Associated with An elevated Risk of Hospital-acquired Acute Kidney Injury

Pinghong He^{1,2*}, Ruixuan Chen^{1*}, Liping Zhou^{1*}, Yanqin Li¹, Licong Su¹, Jin Dong¹, Yan Zha², Yuxin Lin¹, Sheng Nie¹, Fan Fan Hou¹, Xin Xu¹, on behalf of the EACH study investigators

¹Renal Division, Nanfang Hospital, Southern Medical University, The National Clinical Research Center for Kidney Disease, State Key Laboratory of Organ Failure Research, Guangzhou, China;

²Renal Division, Department of Medicine, Guizhou Provincial People's Hospital, Guizhou Provincial Institute of Nephritic and Urinary Disease, Guiyang 550002, Guizhou, China.

*Equal contributors

Correspondence to: Xin Xu, Renal Division, Nanfang Hospital, Southern Medical University, 1838 North Guangzhou Avenue, Guangzhou 510515, China.

Email: xux007@163.com;

Tel: +86-20-61641591; **Fax:** 86-20-87281713.

Abstract word count: 302

Text word count: 2250

Number of References: 29

Number of Tables and Figures: 5

Supplementary Table number: 1

Supplementary Figure number: 1

Supplemental Materials

Table S1. Spearman correlation coefficients among the daily averages of different air pollutants

Pollutant	PM_{2.5}	PM₁₀	NO₂	SO₂	CO	O₃
PM_{2.5}	1.00	0.91	0.69	0.67	0.69	0.09
PM₁₀		1.00	0.68	0.70	0.64	0.12
NO₂			1.00	0.60	0.62	-0.01
SO₂				1.00	0.54	0.08
CO					1.00	-0.18
O₃						1.00




Figure S1. The Flowchart of Patient Selection.