

Supplementary Table 2: Primary and secondary antibodies used in the study.

<i>Primary antibodies</i>			
Antibodies	Dilution for WB	Dilution for IF	Sources
Mouse anti amyloid peptide (clone W02)	/	1:100	Millipore
Rabbit anti GFAP	1:1000	1:100	Cell signaling
Goat anti IBA-1	1:500	1:100	Abcam
Mouse anti β -actin	1:10000	/	Sigma Aldrich
<i>Secondary antibodies</i>			
Antibodies	Dilution for WB	Dilution for IF	Sources
Donkey anti-mouse-Alexa 488	/	1:50	Jackson immunoresearch
Donkey anti-rabbit-RRX	/	1:50	Jackson immunoresearch
Donkey anti-goat RRX	/	1:50	Jackson immunoresearch
Donkey anti goat IgG-HRP	1:1000	/	Santa cruz
Goat anti rabbit IgG-HRP	1:1000	/	Cell Signaling Technology
Horse anti mouse IgG-HRP	1:1000	/	Cell Signaling Technology
<i>Primary antibodies</i>			
Antibodies	Dilution for WB	Dilution for IF	Sources
Mouse anti amyloid peptide (clone W02)	/	1:100	Millipore
Rabbit anti GFAP	1:1000	1:100	Cell signaling
Goat anti IBA-1	1:500	1:100	Abcam
Mouse anti β -actin	1:10000	/	Sigma Aldrich
<i>Secondary antibodies</i>			
Antibodies	Dilution for WB	Dilution for IF	Sources
Donkey anti-mouse-Alexa 488	/	1:50	Jackson immunoresearch
Donkey anti-rabbit-RRX	/	1:50	Jackson immunoresearch
Donkey anti-goat RRX	/	1:50	Jackson immunoresearch
Donkey anti goat IgG-HRP	1:1000	/	Santa cruz
Goat anti rabbit IgG-HRP	1:1000	/	Cell Signaling Technology
Horse anti mouse IgG-HRP	1:1000	/	Cell Signaling Technology