

Supplementary table 2. Echocardiographic evaluation during first 24 hours

Measurements	n = 38
Left ventricular posterior wall thickness, mm, mean (SD)	11 (2)
Septal thickness, mm, mean (SD)	11 (2)
Left ventricular end-diastolic dimension, mm, mean (SD)	50 (10)
Left ventricular end-systolic dimension, mm, mean (SD)	37 (12)
Left ventricular ejection fraction, n (%)	
<20 %	2 (5)
20 – <30 %	3 (8)
30 – <40 %	7 (18)
40 – <50 %	13 (35)
50 – <60 %	11 (29)
≥60 %	2 (5)
Left ventricular fractional shortening, %, mean (SD)	27 (10)
Wall motion score index, score, median (Q1–Q3)	1.5 (1.1–1.8)
Left ventricular outflow tract velocity time integral, cm, mean (SD)	16.6 (4.3)
Right ventricular basal diameter, mm, mean (SD)	39.4 (6.4)
Tricuspid annular plane systolic excursion, mm, mean (SD)	18.5 (4.3)
ST-elevated myocardial infarction	21 (55)
Non-ST-elevated myocardial infarction	1 (2)
Old myocardial infarction, n (%)	6 (16)
Valvular pathology ^{1,2}	13 (34)

¹ Excluding mild tricuspid valve insufficiency.

² One mild aortic valve stenosis; one moderate aortic valve insufficiency; six mild and one moderate mitral valve insufficiency; three moderate and one severe tricuspid valve insufficiency.

SD: standard deviation; Q1–Q3: first to third quartile.