

S2 Table: Treatment received by the patients in COVID-19 wave 1 and wave 2

Wave 1	Acute phase (n=25)				Recovery phase (n=19)			
	Anti-infectives	Hydroxy-chloroquin	Steroids	Other Immuno-suppression	Anti-infectives	Hydroxy-chloroquin	Steroids	Other Immuno-suppression
Total	24/25 (96%)	11/25 (44%)	7/25 (28%)	4/25 (16%)	12/19 (63%)	0/20 (0%)	7/19 (36%)	3/19 (16%)

Wave 2	Acute phase (n=38)				Recovery phase (n=21)			
	Anti-infectives	Hydroxy-chloroquin	Steroids	Other Immuno-suppression	Anti-infectives	Hydroxy-chloroquin	Steroids	Other Immuno-suppression
Total	37/38 (97%)	0/38 (0%)	38/38 (100%)	3/38 (7.9%)	9/21 (42%)	0/21 (0%)	10/21 (47.6%)	1/21 (4.7%)

Definitions:

Anti-infective: Use of either/or antibiotics, antivirals, antifungals in the last 48 hours before sampling. The use of prophylactic anti-infective treatment (e.g. trimethoprim-sulfamethoxazole 3x/week) was not considered in this analysis.

Hydroxychloroquin: Use of hydroxychloroquin in the last 7 days before sampling

Steroids: Any use of steroids equivalent to 20mg prednisone per day in the last 48 hours before sampling