

Supplementary Table 2. Primers used for PCR amplifications

Gene	Primer (5'-3')	
CPVL	Forward	TCAACCTGAACGGAATTGCTA
	Reverse	GAAGGATCACTTGTTAAGTCGC
IRF9	Forward	CCGAAACTCCGGAAGTGGG
	Reverse	GGCCTTGAAGAAGGCAGCAT
NF- κ B	Forward	CGGCTGAGTCCTGCTCCTTCC
	Reverse	TGGAGGCTGCCTGGATCACTTC
IFITM2	Forward	AGACCTCCGTGCCTGACCATG
	Reverse	CGTCGCCAACCATCTTCCTGTC
IRF1	Forward	CACTGTCGCCATGTGCTGTCAG
	Reverse	TGCCACTCCGACTGCTCCAAG
BAK	Forward	GGACGACATCAACCGACGCTATG
	Reverse	AACAGGCTGGTGGCAATCTTGG
TAP1	Forward	ACAGCCAGTTACAGGTGGAGCAG
	Reverse	CCGGATAGCGCCTCCTTCCAG
IFITM3	Forward	GCATTGCCTACTCCGTGAAGTC
	Reverse	ACTTGGCGGTGGAGGCATAGG
IFITM1	Forward	GATCAACATCCACAGCGAGACCTC
	Reverse	GACTTCACGGAGTAGGCGAATGC
PSMB8	Forward	GAGACAGCTATTCTGGAGGCGTTG
	Reverse	GTGCAGCAGGTCCTGACATCTG
IFI35	Forward	CTCGGTACTGGTGCTCAACATTCC
	Reverse	ACTCAGAGGTGAAGACTGCTAGGC
BCl-2	Forward	ACTGAGTACCTGAACCGGCATC
	Reverse	GGAGAAATCAAACAGAGGTTCGC
BAX	Forward	AGTGTCTCAGGCGAATTGGC
	Reverse	CACGGAAGAAGACCTCTCGG
BTK	Forward	CCAATGGCTGCCTCCTGAACTAC
	Reverse	TCGGTGAAGGAACTGCTTTGACTC
GAPDH	Forward	TGACTTCAACAGCGACACCCA
	Reverse	CACCCTGTTGCTGTAGCCAAA