nature portfolio

Corresponding author(s):	Stephan Riesenberg
Last updated by author(s):	Jan 5, 2022

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our Editorial Policies and the Editorial Policy Checklist.

~					
5	tа	ŤΙ	101	h	2

For	all statistical a	nalyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.				
n/a	Confirmed					
	🗶 The exac	e exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement				
	🗶 A statem	statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly				
×		The statistical test(s) used AND whether they are one- or two-sided Only common tests should be described solely by name; describe more complex techniques in the Methods section.				
×	A description of all covariates tested					
×	A descrip	A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons				
	A full des AND vari	A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)				
x	For null hypothesis testing, the test statistic (e.g. <i>F</i> , <i>t</i> , <i>r</i>) with confidence intervals, effect sizes, degrees of freedom and <i>P</i> value noted <i>Give P values as exact values whenever suitable.</i>					
×	For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings					
×	For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes					
×	Estimates of effect sizes (e.g. Cohen's <i>d</i> , Pearson's <i>r</i>), indicating how they were calculated					
Our web collection on <u>statistics for biologists</u> contains articles on many of the points above.						
Software and code						
Policy information about <u>availability of computer code</u>						
Da	ata collection No software was used.					
Da	CRISPResso (v1), SAMtools (v1.12), Integrative Genomics Viewer (v2.10), GraphPad Prism (v6)					
For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio guidelines for submitting code & software for further information						

Data

Policy information about availability of data

All manuscripts must include a <u>data availability statement</u>. This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our policy

Source Data are provided with this paper. The NGS data generated in this study have been deposited in the Dryad database under accession code dryad.2rbnzs7mk (https://datadryad.org/stash/dataset/doi:10.5061/dryad.jwstqjqb7). Data are also available on request from authors.

Field-specific reporting					
Please select the or	ne below that is	the best fit for your research. If you are not sure, read the appropriate sections before making your selection.			
X Life sciences	□ В	ehavioural & social sciences			
For a reference copy of t	the document with a	Ill sections, see <u>nature.com/documents/nr-reporting-summary-flat.pdf</u>			
		ıdy design			
		points even when the disclosure is negative.			
Sample size	At least two independent biological replicates for each condition. This was sufficient as the editing efficiency difference between replicates of a respective group was very small compared to the strong difference observed between the compared groups.				
Data exclusions	No data were ex	ata were exluded from analysis.			
Replication	All attempts of replication (n = 2 or 3) were successful.				
Randomization	No randomization	randomization was necessary for this study because experiments were done under well controlled identical conditions.			
Blinding	Samples were prepared unblinded but in parallel using cells grown under identical conditions. Analysis was performed based on numerical sample names, without the identity of the samples known during the analysis.				
We require information	on from authors a	pecific materials, systems and methods bout some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.			
Materials & exp	perimental sy	ystems Methods			
- 					
Antibodies					
■ Eukaryotic cell lines ■ Flow cytometry					
Palaeontology and archaeology MRI-based neuroimaging					
Animals and other organisms					
Human research participants Clinical data					
Dual use research of concern					
Eukaryotic c	ell lines				
Policy information a	about <u>cell lines</u>				
Cell line source(s) 409-B2 hiPSCs (Riken BioResource Center), HEK293 cells (ECACC, 85120602), K562 cells (ECACC, 89121407),		409-B2 hiPSCs (Riken BioResource Center), HEK293 cells (ECACC, 85120602), K562 cells (ECACC, 89121407), Jurkat cells			

Cell line source(s)

409-B2 hiPSCs (Riken BioResource Center), HEK293 cells (ECACC, 85120602), K562 cells (ECACC, 89121407), Jurkat cells (ATCC, TIB-152), Caco-2 cells (DSMZ, ACC169), SNL 76/7 (ECACC, 07032801), CHO-K1 cells (ATCC, CCL61)

Authentication

All cell lines were authenticated by the supplier via certificate of analysis and additionally in-house by checking morphology.

Mycoplasma contamination

Commonly misidentified lines (See ICLAC register)

No commonly misidentified cell lines were used in the study.